

Measuring long-term outcomes after injury: current issues and future directions

Ali Salim,¹ Deborah M Stein ², Ben L Zarzaur,³ David H Livingston ⁴

¹Surgery, Brigham and Women's Hospital - Harvard Medical School, Boston, Massachusetts, USA

²Surgery, University of Maryland School of Medicine, Baltimore, Maryland, USA

³Surgery, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA

⁴Surgery, Rutgers-New Jersey Medical School, Newark, New Jersey, USA

Correspondence to

Dr David H Livingston; livingst@rutgers.edu

Received 29 November 2022

Accepted 16 February 2023

SUMMARY

Maximizing long-term outcomes for patients following injury is the next challenge in the delivery of patient-centered trauma care. The following review outlines three important components in trauma outcomes: (1) data gathering and monitoring, (2) the impact of traumatic brain injury, and (3) trajectories in recovery and identifies knowledge gaps and areas for needed future research.

For the past 50 years, improvements in the delivery of trauma center care have markedly decreased mortality after injury. Unfortunately, our knowledge of what happens to survivors once they leave the hospital is limited. The reasons for this are multifactorial. For many years, 'discharged alive' was sufficient and in-hospital mortality was the metric on which trauma center care was judged. There was also little interest or regulatory requirement to collect data beyond discharge.

There has been increased recognition that inpatient trauma care makes up only a very small, although vital, part of a patient's recovery. The focus on long-term outcomes is necessary to begin to understand the degree of disability our patients experience and how these outcomes may be modulated following discharge to optimize recovery.^{1,2} In 2021, at the annual meeting of the American Association of the Surgery of Trauma, a panel of experts provided an overview of some of the important issues that need to be addressed when considering long-term outcomes following injury. The following paper outlines three important components in trauma outcomes: data gathering and monitoring, the impact of traumatic brain injury, and trajectories in recovery. More importantly, it begins to identify those gaps in our knowledge and those areas which require further investigation and study. The overall focus on these outcomes must become the next frontier in overall patient-centered trauma care. It is hoped that these types of future investigations will ultimately improve our patients' quality of life (QoL) and reintegration back to family and society similar to the impact past studies had on improving overall inpatient survival after injury.

DATA GATHERING AND MONITORING

Data are the key to understanding and improving patient care. This concept began with the mandatory requirement for all trauma centers to have individual registries and has further expanded into the robust and important Trauma Quality Improvement Project. However, these data comprise prehospital

and in-hospital events with minimal collection of postdischarge data.

One of the longest and most successful endeavors in tracking long-term trauma outcomes is the Victoria State Trauma Registry (VSTR) in Australia; under the direction of Gabbe *et al.*³⁻⁷ In addition to the standard inpatient trauma registry data, the VSTR re-engages patients at 6, 12, and 24 months after injury to collect health-related QoL, function, disability, pain, and return to work outcomes. These longitudinal data on physical and behavioral health, social, and cognitive impairments uncovered by this population-based registry have provided tremendous insights into the outcome of trauma patients in that province. One of the key findings from this work demonstrates that there is significant impairment in all domains which does not substantially improve for the 3 years postdischarge.⁸ These data paint a rather bleak picture of life following major trauma and provide an impetus to formulate interventions that can positively impact the recovery of trauma patients. It is these data gathering models which will be able to assess the impact and potential successes of these interventions.

In the USA, two similar longitudinal studies have been launched over the past 20 years including the Functional Outcomes and Recovery after Trauma (FORTE).⁹⁻¹¹ Similar to the VSTR, the FORTE project is a collaboration between three level 1 trauma centers in Boston to determine the feasibility, sustainability, and value of the routine collection of long-term outcomes after injury. With data collected at 6 and 12 months postinjury from over 4500 patients, the FORTE project is one of the largest repositories for long-term trauma outcomes in the USA. Some of the key observations suggest and reiterate the findings of others, that patient factors (age, gender, race, education) rather than injury-related factors (injury severity or location) are more predictive of long-term outcomes¹²⁻¹⁵ and that poor long-term outcomes tend to be associated with each other within and across different health domains (behavioral health (eg, post-traumatic stress disorder (PTSD)), physical, social).^{16,17} Potentially modifiable factors such as social support were found to potentially influence the recovery trajectories of these patients.^{18,19}

The impetus and momentum for US trauma centers to collect long-term outcomes has been slow but there has been a small increase in the number of published research articles over the last several years. Unfortunately, there is considerable variability in the metrics used to measure outcomes. In 2019, a consensus conference of key stakeholders was convened to propose the optimal metrics to assess

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Salim A, Stein DM, Zarzaur BL, *et al.* *Trauma Surg Acute Care Open* 2023;**8**:e001068.

long-term outcomes as well as the appropriate instruments with which to measure them.²⁰ Subsequently, the National Trauma Research Action Plan collaborative has proposed 14 patient-reported outcome measures (PROMs) across 13 core domains including activities of daily living, alcohol and substance use, behavioral health metrics (such as PTSD and depression), and measures of general and trauma-specific QoL.²¹ One substantial concern is that these quantitative metrics, while useful for many surgical conditions may fail to capture the true picture and nuances of outcomes in trauma patients. More importantly, they do not provide insight into potential solutions to modulate outcomes. A recent descriptive study by Rosenberg *et al* demonstrated that recovery of both the physical body and the ‘self’, a composite of one’s roles, values, identities, and beliefs was essential to achieving favorable subjective outcomes.²² These complex qualitative findings which may not be reflective of quantitative ordinal scales in the same patients have been suggested by others²³ and it is likely that to obtain a true assessment of long-term outcomes following trauma will require both qualitative and quantitative measures.

Unfortunately, obtaining high-quality data are both costly and labor intensive. Interest has recently emerged in other fields in measuring long-term outcomes through indirect measures such as digital phenotyping; a technique of passively collecting data through an individual’s smartphone sensors and use.^{24–26} This methodology offers the possibility of obtaining objective data without the manpower needed for telephone surveys. For example, the collection of global positioning system data might possibly be used to study physical mobility and community integration after trauma based on examining how a person spends their time at different locations.

Determining the best methodology to gather these data is an area of ongoing research and will need to meet the aims of long-term outcome data collection: (1) low cost, (2) minimal burden to patients and data registrars, and (3) production of sufficiently granular and high-quality data that can be used to measure outcomes that matter. The best strategy may in fact be a combination that deploys surveys and tasks via email, text, and apps with phone interviews to collect PROMs data. Recently, the VSTR has demonstrated that on-line report was almost as good as phone interviews.²⁷ However, as outlined above, the issue of human contact and obtaining more qualitative information may ultimately be necessary in truly assessing recovery.

TRAUMATIC BRAIN INJURY

Traumatic brain injury (TBI) remains a huge public health problem which accounts for 80 000 permanently disabled individuals each year in the USA.²⁸ Additionally, TBI disproportionately affects certain groups that suffer from healthcare disparities such as racial and ethnic minorities and people living in rural areas.^{29–33} The impact of TBI on overall trauma outcomes cannot be underestimated. In a retrospective study of severe trauma patients by Livingston *et al*, those patients with a TBI did substantially worse at 3.3 years than those free of brain injury.³⁴ In addition, survivors of TBI are at an increased risk of depression and PTSD, suicide, and substance abuse.

There is a well-described natural history of an acute severe TBI, but the ultimate recovery and outcome of an individual patient is exceedingly variable which can appear random to the lay public and even to medical personnel. Determining the recovery of a patient following moderate-to-severe TBI is difficult and fraught with uncertainty.³⁵ Prognostic models have been created such as ‘Crash’ and ‘Impact TBI’ that are able to provide

reasonable predictions for unfavorable outcome at 2 weeks and/or 6 months and death based on clinical and radiographic variables.^{36,37} One systematic review of 27 studies demonstrated that early imaging and evaluation of the deep cerebral structures can yield significant prognostic information.³⁸ It is advisable to be extremely cautious as these tools only provide early and somewhat gross prognostic information.

There is no question that many patients with moderate-to-severe TBI make great strides in the first year following injury.³⁹ Unfortunately, TBI is a chronic disease with the best data on longitudinal follow-up coming from the TBI models systems from the National Institute on Disability and Rehabilitation Research. Unfortunately, the data from 1 to 5 years do not paint a promising picture for long-term improvement.⁴⁰ In these data, only 25% of patients had any improvement over time. For the remaining 75%, one-third die and one-third suffer marked deterioration.

Even if we were able to prognosticate with some measure of certainty following TBI, defining a ‘favorable’ outcome following a TBI is subjective and dependent on the outcome scales used. In many studies, the Glasgow Outcome Scale (GOS) or GOS-extended version is used to define a ‘favorable’ outcome. However, these scales have an exceedingly low threshold for what is considered ‘favorable’ to an individual patient or family. For example, being unable to shop without assistance, not being able to return to work, and severe restrictions in social and leisure activities are all considered ‘favorable’ outcome by these scales but would not likely be favorable by most people.

Patients who sustain a moderate or severe TBI have a reduced life expectancy by 9 years. Survivors alive 5 years after a TBI suffer a litany of ongoing medical and social problems; 57% are moderately or severely disabled, 55% who were previously employed are unemployed, 50% require at least one hospital readmission, 33% are dependent for their activities of daily living, 29% are not ‘satisfied’ with life, 29% have substance use disorders, and 12% reside in chronic care facilities.^{41–43} Physical symptoms, often lifelong, such as headache, dizziness and nausea, fatigue, and lethargy are common as well as cognitive deficits such as disturbances of attention and memory, slow cognition, inability to multitask, increased distractibility, and impaired executive function.⁴⁴ Memory failure is the most commonly reported cognitive issue following TBI, while mental slowness and attention disorders are the next most common occurring in 57% and 52% of patients.^{45–47} Fatigue is extremely common and is reported by 30%–70% of patients.^{48,49} Additionally, ‘anosognia’ or lack of awareness of cognitive and behavioral disorders occurs in 76%–97% of patients following severe TBI, which contributes to difficulty in treatment and poor outcomes.^{47,50–52} Behavioral disturbances such as personality changes, depression, anxiety, impulsivity, irritability, emotional lability, apathy, and depression occur frequently as well.⁴⁴ In one study, 2 years after TBI, irritability was found in 67% of patients, a lack of initiative in 44%, and socially inappropriate behavior in 26%.⁵² Nearly 60% of patients experienced personality change after an acquired brain injury which is associated with a high subjective burden for the patients’ caregivers.^{52,53}

Therefore, we must ask ourselves whether any of these outcomes are currently modifiable. Is a poor outcome or a ‘favorable’ outcome with severe personality and cognitive disturbances a forgone conclusion following TBI? Without long-term longitudinal data, these answers will never be known. We believe that there may be opportunities both in inpatient and outpatient treatments that may affect outcomes. At present, social factors such as educational attainment⁵⁴ and preinjury employment⁵⁵

seem to be the best predictors of functional recovery after controlling for severity of injury. This implies that social factors are potentially modifiable factors that could be leveraged to improve outcomes for patients with severe TBI in the future. These data clearly suggest that overall impact and trajectory of recovery for patients with and without TBI are very different. In contrast to other trauma studies which mix patients with and without TBI, we would strongly suggest that studies examining long-term outcomes following injury should separate these populations.

TRAJECTORIES AFTER INJURY

It is increasingly clear that recovery after injury is a complex and dynamic process with many factors that influence how fully an individual patient may recover. To better understand recovery and to design interventions that may be effective in improving long-term outcomes, it is imperative that we reconsider the underlying assumptions that we have regarding recovery. The most common assumption as described by Iwashyna is that most patients follow a ‘Big Hit’ trajectory following injury.⁵⁶ In this model, patients have a severe and sudden decline in function followed by a slow recovery over several months until reaching a new baseline. But data from several studies indicate that this may not be the prevailing or the only trajectory followed during recovery.

Until recently, most of the statistical methods used to analyze retrospective and prospectively collected long-term outcomes force all the data used to fit a model in which a mean curve is constructed with the variance of the curve made up of random and systemic error. This approach to analysis forces outliers to fit the underlying shape of the outcome curve which could obscure the existence of other recovery trajectories. One way to overcome this problem is to use group-based trajectory modeling (GBTM). GBTM is a type of finite mixture modeling that does not make assumptions about the number or shape of trajectories that may exist in longitudinal data. When applied to longitudinally collected data among injured people, several possible trajectories of recovery following injury emerge (figure 1).⁵⁷

Using the data from the Health and Retirement Study, which surveys a cohort of people 50 and older every 2 years, Bell *et al* applied GBTM to a measure of functional limitation 2 years before and 8 years after a reported injury.⁵⁸ The analysis demonstrated that five different trajectories existed in the longitudinal functional limitation data. Only 46% of the cohort followed the

‘Big Hit’ trajectory. Three other trajectories were identified that changed over time: the ‘slow burn’, the ‘long-term improvement’, and the ‘low functioning’. The ‘slow burn’ trajectory was characterized by an initial steep decline in function followed by a less severe, but still declining trajectory. The ‘long-term improvement’ trajectory improved slowly after an initial decrease in functional status. Finally, the ‘low functioning’ trajectory was characterized by low functioning at the time of injury and suffered a decrease in function but improved over time.

While these data demonstrate the existence of different recovery trajectories, there was no information available regarding the type or severity of injury that the people in the Health and Retirement Survey cohort suffered. In the Socio-economic Status and Post-Injury Quality of Life and Functional Ability Study, 500 non-TBI or spinal cord injury survivors were followed for 1 year. GBTM was used to identify classes of individuals within the study sample that had varying and distinct recovery patterns over 12 months.⁵⁹ This analysis used the Short Form-36 as the outcome measures for functional and psychological outcomes and identified three physical and five behavioral health trajectories. For the physical outcomes, only 24% of patients followed the expected recovery trajectory which has a sharp decline followed by rapid recovery. The remainder followed worse trajectories. For the behavioral health outcomes, a similar pattern was observed. Only 23% followed the expected trajectory with one-third of patients following a resilient recovery

Box 1 Challenges and future directions for optimizing trauma outcomes

Data collection

1. How best to capture high-quality data at the lowest possible cost?
2. What standardized and validated screening tools should be used to detect changes in psychological status (eg, post-traumatic stress disorder (PTSD)), physical function, caregiver burden, sleep disturbances, and perceived social support?
3. How to incorporate qualitative data into a measure of trauma outcomes?
4. How long should trauma patients be followed to determine ‘final outcome measures’?

Traumatic brain injury

1. The recognition that the outcomes of patients with traumatic brain injury (TBI) are so different that they should not be mixed with non-TBI patients.
2. Identify and define favorable outcomes in TBI.
3. Can the TBI outcomes be modified during inpatient care?
4. Can TBI outcomes be modified and improved with targeted outpatient care?

Trajectory and interventions

1. Identification of which groups may benefit the most from post-trauma center discharge intervention?
2. How often should behavioral health (eg, PTSD and depression) screening be done?
3. What is the optimal composition of the outpatient team to provide comprehensive collaborative care?
4. How does discharge location affect trajectory and outcomes?
5. What is the effect and impact of caregivers on trauma outcomes?
6. Does caregiver support improve outcomes?
7. What is the effect of peer support programs on trauma outcomes?




Figure 1 Various trajectory models for trauma recovery.

trajectory in which no decrease in behavioral health outcomes was observed. In summary, in this vulnerable population, over 75% of the non-neurological injury survivors followed suboptimal physical recovery trajectories and nearly 50% suffered a long-term decrease in psychological recovery. This was true even after accounting for differences in injury severity. The important underlying question in these observational database studies is ‘Can these poor trajectories be modified through intervention?’

Identifying those who might benefit from an intervention to improve outcomes is problematic. Similar to other medical interventional studies there exist three populations: those who are destined to improve, and which interventions will have no harm but may offer no benefit, those who are destined to do worse and which interventions are doomed to fail, and the target group where the interventions may improve outcomes. It is also important to recognize that not all outcomes may be modifiable in the same way. In terms of outcome trajectories after injury, we would like to be able to identify the last group so that we can have the most impact. Unfortunately, at the present time, no good screening criteria exist to indicate which patients will follow one of the dynamic recovery trajectories following injury. However, data are emerging that indicate certain groups may be at higher risk of suffering a poor outcome trajectory than others. Older patients, women, and those with multiple medical comorbidities are at risk for following one of the less desirable outcome trajectories.⁵⁹ Pre-existing psychological comorbidities and substance abuse are also an indication of poor functional and psychological outcomes.^{17 60 61} There is mounting evidence that poor perceived social support is associated with following a suboptimal outcome trajectory.^{19 62 63}

Taken together, the data from cross-sectional and longitudinal outcome studies indicate that there are persistent impairments of physical and psychological outcomes for patients after injury.^{7 9 34 58 59 64} While some patients follow the ‘Big Hit’ outcome trajectory, more have a very different recovery experience. Also, biopsychosocial impairment persists for many months after injury in a time where patients are in an outpatient setting and not readily available for screening and intervention. Injury survivors are also at risk from fragmentation of care which is a major factor in poor outcomes in patients with complex biopsychosocial needs. More worrisome are those who follow one of the trajectories that are initially high functioning and then suffer some setback weeks after their injury that leads to a worsening of their overall QoL. Ultimately, recovery from injury is a dynamic process that occurs over many weeks to months with most people reaching a steady state at some point 1 year after injury.

There are evidence-based interventions that exist from which we can learn about how to improve long-term outcomes after injury. Screening for post-traumatic stress in the inpatient setting following injury and following a stepped care pathway that includes prescribing medications reduces the risk of the development of PTSD and depression in the months after injury.^{65 66} Despite limitations, this type of screening and treatment should be a part of any comprehensive program to improve the overall QoL of injured patients.

To augment inpatient focused interventions, we need to consider the fact that recovery takes many months to years, that it is a dynamic process, and that care fragmentation and social isolation are significant threats to achieving ideal outcomes after injury. To address these threats, new and novel approaches to the outpatient management of injury survivors must be considered. Furthermore, trauma surgeons and trauma centers are in an ideal position to implement programs for the long-term care of trauma patients. A type of intervention that may hold promise

is a collaborative care intervention. Collaborative care interventions are designed to meet the needs of biopsychosocially complex patients who are at particular risk from fragmented care in the outpatient setting. There have been several randomized controlled trials that demonstrate the effectiveness of improving outcomes or delaying decline in patients with dementia, multiple comorbidities, social frailty, and in the recovery from critical illness.^{67–69} The collaborative care delivery model is a multidisciplinary process to standardize and streamline care for selected case types—particularly for those with complex cognitive, physical, or psychological morbidity. Care coordination includes these features: (1) a patient-centered approach focused on the goals of the patient; (2) team and measurement-based care plan with each team member having the appropriate skills and resources to carry out their role; (3) a process to track the efficacy of a personalized care plan to meet the complex cognitive, physical, and psychological morbidity of each patient longitudinally to inform care decisions and to facilitate communication and coordination of care across team members and across sites of care; (4) a stepped care approach that provides the patient with the dose and duration of evidence-based treatment necessary to meet their goals of care; and (5) access to specialty care or community resources for those patients whose goals are not being met with current interventions. There are no clear data regarding how frequently follow-up should occur nor how long it should continue but all should recognize that serious injury is a life-changing event.^{70–72}

CONCLUSION

We believe that maximizing long-term outcomes is the next challenge that trauma centers must face. The data presented (box 1) attempts to outline both the current state of knowledge and areas where future study is required.

Contributors All authors participated in the construction and editing of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Deborah M Stein <http://orcid.org/0000-0003-3683-3963>

David H Livingston <http://orcid.org/0000-0003-0427-0393>

REFERENCES

- Gissane W, Bull J, Roberts B. Sequelae of road injuries: a review of one year's admission to an accident hospital. *Injury* 1970;1:195–203.
- Richmond TS, Kauder D, Schwab CW. A prospective study of predictors of disability at 3 months after non-central nervous system trauma. *J Trauma* 1998;44:635–42.
- Gabbe BJ, Cameron PA, Hannaford AP, Sutherland AM, McNeil JJ. Routine follow up of major trauma patients from trauma registries: what are the outcomes? *J Trauma* 2006;61:1393–9.
- Gabbe BJ, Simpson PM, Sutherland AM, Wolfe R, Lyons RA, Cameron PA. Evaluating time points for measuring recovery after major trauma in adults. *Ann Surg* 2013;257:166–72.
- Gabbe BJ, Sutherland AM, Hart MJ, Cameron PA. Population-based capture of long-term functional and quality of life outcomes after major trauma: the experiences of the Victorian state trauma registry. *J Trauma* 2010;69:532–6.

- 6 Gabbe BJ, Simpson PM, Sutherland AM, Palmer CS, Williamson OD, Butt W, Bevan C, Cameron PA. Functional and health-related quality of life outcomes after pediatric trauma. *J Trauma* 2011;70:1532–8.
- 7 Gabbe BJ, Simpson PM, Harrison JE, Lyons RA, Ameratunga S, Ponsford J, Fitzgerald M, Judson R, Collie A, Cameron PA. Return to work and functional outcomes after major trauma: who recovers, when, and how well? *Ann Surg* 2016;263:623–32.
- 8 Gabbe BJ, Simpson PM, Cameron PA, Ponsford J, Lyons RA, Collie A, Fitzgerald M, Judson R, Teague WJ, Braaf S, *et al.* Long-term health status and trajectories of seriously injured patients: a population-based longitudinal study. *PLoS Med* 2017;14:e1002322.
- 9 Haider AH, Herrera-Escobar JP, Al Rafai SS, Harlow AF, Apoj M, Nehra D, Kasotakis G, Brasel K, Kaafarani HMA, Velmahos G, *et al.* Factors associated with long-term outcomes after injury: results of the functional outcomes and recovery after trauma emergencies (Forte) multicenter cohort study. *Ann Surg* 2020;271:1165–73.
- 10 Rios-Diaz AJ, Herrera-Escobar JP, Lilley EJ, Appelton JR, Gabbe B, Brasel K, deRoon-Cassini T, Schneider EB, Kasotakis G, Kaafarani H, *et al.* Routine inclusion of long-term functional and patient-reported outcomes into trauma registries: the Forte project. *J Trauma Acute Care Surg* 2017;83:97–104.
- 11 Herrera-Escobar JP, Seshadri AJ, Rivero R, Toppo A, Al Rafai SS, Scott JW, Havens JM, Velmahos G, Kasotakis G, Salim A, *et al.* Lower education and income predict worse long-term outcomes after injury. *J Trauma Acute Care Surg* 2019;87:104–10.
- 12 Herrera-Escobar JP, El Moheb M, Ranjit A, Weed C, Brasel K, Kasotakis G, Kaafarani HMA, Velmahos G, Nehra D, Haider AH, *et al.* Sex differences in long-term outcomes after traumatic injury: a mediation analysis. *Am J Surg* 2021;222:842–8.
- 13 Chun Fat S, Herrera-Escobar JP, Seshadri AJ, Al Rafai SS, Hashmi ZG, de Jager E, Velmahos C, Kasotakis G, Velmahos G, Salim A, *et al.* Racial disparities in post-discharge healthcare utilization after trauma. *Am J Surg* 2019;218:842–6.
- 14 Herrera-Escobar JP, Rivero R, Apoj M, Geada A, Villanyi M, Blake D, Nehra D, Velmahos G, Kaafarani HMA, Salim A, *et al.* Long-term social dysfunction after trauma: what is the prevalence, risk factors, and associated outcomes? *Surgery* 2019;166:392–7.
- 15 Jurkovich G, Mock C, MacKenzie E, Burgess A, Cushing B, deLateur B, McAndrew M, Morris J, Swiontkowski M. The sickness impact profile as a tool to evaluate functional outcome in trauma patients. *J Trauma* 1995;39:625–31.
- 16 Herrera-Escobar JP, Apoj M, Weed C, Harlow AF, Al Rafai SS, Lilley E, Kasotakis G, Brasel K, Kaafarani HMA, Velmahos G, *et al.* Association of pain after trauma with long-term functional and mental health outcomes. *J Trauma Acute Care Surg* 2018;85:773–9.
- 17 Herrera-Escobar JP, Seshadri AJ, Stanek E, Lu K, Han K, Sanchez S, Kaafarani HMA, Salim A, Levy-Carrick NC, Nehra D. Mental health burden after injury: it's about more than just posttraumatic stress disorder. *Ann Surg* 2021;274:e1162–9.
- 18 Nehra D, Herrera-Escobar JP, Al Rafai SS, Havens J, Askari R, Nitzschke S, Velmahos G, Kasotakis G, Brasel KJ, Levy-Carrick N, *et al.* Resilience and long-term outcomes after trauma: an opportunity for early intervention? *J Trauma Acute Care Surg* 2019;87:782–9.
- 19 Orlas CP, Herrera-Escobar JP, Hau KM, Velmahos A, Patel N, Sanchez S, Kaafarani HMA, Salim A, Nehra D. Perceived social support is strongly associated with recovery after injury. *J Trauma Acute Care Surg* 2021;91:552–8.
- 20 Sakran JV, Ezzeddine H, Schwab CW, Bonne S, Brasel KJ, Burd RS, Cuschieri J, Ficke J, Gaines BA, Giacino JT, *et al.* Proceedings from the consensus conference on trauma patient-reported outcome measures. *J Am Coll Surg* 2020;230:819–35.
- 21 Herrera-Escobar JP, Price MA, Reidy E, Bixby PJ, Hau K, Bulger EM, Haider AH, National Trauma Research Action Plan (NTRAP) Long-term Patient-Reported Outcomes Panel. Core outcome measures for research in traumatic injury survivors: the national trauma research action plan modified delphi consensus study. *J Trauma Acute Care Surg* 2022;92:916–23.
- 22 Rosenberg G, Zion SR, Shearer E, Berekyei Merrell S, Abadilla N, Spain DA, Crum AJ, Weiser TG. What constitutes a “successful” recovery? Patient perceptions of the recovery process after a traumatic injury. *Trauma Surg Acute Care Open* 2020;5:e000427.
- 23 Ekegren CL, Braaf S, Ameratunga S, Ponsford J, Nunn A, Cameron P, Lyons RA, Gabbe BJ. Adaptation, self-motivation and support services are key to physical activity participation three to five years after major trauma: a qualitative study. *J Physiother* 2020;66:188–95.
- 24 Cote DJ, Barnett I, Onnala J-P, Smith TR. Digital phenotyping in patients with spine disease: a novel approach to quantifying mobility and quality of life. *World Neurosurg* 2019;126:e241–9.
- 25 Panda N, Solsky I, Huang EJ, Lipsitz S, Pradarelli JC, Delisle M, Cusack JC, Gadd MA, Lubitz CC, Mullen JT, *et al.* Using smartphones to capture novel recovery metrics after cancer surgery. *JAMA Surg* 2020;155:123–9.
- 26 Wright AA, Raman N, Staples P, Schonholz S, Cronin A, Carlson K, Keating NL, Onnala J-P. The hope pilot study: harnessing patient-reported outcomes and biometric data to enhance cancer care. *JCO Clin Cancer Inform* 2018;2:1–12.
- 27 Gabbe BJ, Hart MJ, Brown A, McLellan S, Morgan MJ, Beck B, de Steiger RS, Cameron PA. Does patient preference for online or telephone follow-up impact on response rates and data completeness following injury? *J Trauma Acute Care Surg* 2022;93:679–85.
- 28 Taylor CA, Bell JM, Breiding MJ, Xu L. Traumatic brain injury-related emergency department visits, hospitalizations, and deaths—United States, 2007 and 2013. *MMWR Surveill Summ* 2017;66:1–16.
- 29 The CDC, NIH, DoD, and VA Leadership Panel. Report to congress on traumatic brain injury in the united states: understanding the public health problem among current and former military personnel. Centers for Disease Control and Prevention (CDC), the National Institutes of Health (NIH), the Department of Defense (DoD), and the Department of Veterans Affairs (VA), 2013.
- 30 Stubbs JL, Thornton AE, Seveck JM, Silverberg ND, Barr AM, Honer WG, Panenka WJ. Traumatic brain injury in homeless and marginally housed individuals: a systematic review and meta-analysis. *Lancet Public Health* 2020;5:e19–32.
- 31 Durand E, Chevignard M, Ruet A, Dereix A, Jourdan C, Pradat-Diehl P. History of traumatic brain injury in prison populations: a systematic review. *Ann Phys Rehabil Med* 2017;60:95–101.
- 32 St Ivany A, Schminkey D. Intimate partner violence and traumatic brain injury: state of the science and next steps. *Fam Community Health* 2016;39:129–37.
- 33 Chapal A. Traumatic brain injury: outcomes of a rural versus urban population over a 5-year period. *Hawaii Med J* 2007;66:318–21.
- 34 Livingston DH, Tripp T, Biggs C, Lavery RF. A fate worse than death? Long-term outcome of trauma patients admitted to the surgical intensive care unit. *J Trauma* 2009;67:341–8.
- 35 Multi-Society Task Force on PVS. Medical aspects of the persistent vegetative state (2). *N Engl J Med* 1994;330:1572–9.
- 36 MRC CRASH Trial Collaborators, Perel P, Arango M, Clayton T, Edwards P, Komolafe E, Pocock S, Roberts I, Shakur H, Steyerberg E, *et al.* Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ* 2008;336:425–9.
- 37 Steyerberg EW, Mushkudiani N, Perel P, Butcher I, Lu J, McHugh GS, Murray GD, Marmarou A, Roberts I, Habbema JDF, *et al.* Predicting outcome after traumatic brain injury: development and international validation of prognostic scores based on admission characteristics. *PLoS Med* 2008;5:e165.
- 38 Haghbayan H, Boutin A, Laflamme M, Lauzier F, Shemilt M, Moore L, Zarychanski R, Douville V, Fergusson D, Turgeon AF. The prognostic value of MRI in moderate and severe traumatic brain injury: a systematic review and meta-analysis. *Crit Care Med* 2017;45:e1280–8.
- 39 McCrea MA, Giacino JT, Barber J, Temkin NR, Nelson LD, Levin HS, Dikmen S, Stein M, Bodien YG, Boase K, *et al.* Functional outcomes over the first year after moderate to severe traumatic brain injury in the prospective, longitudinal TRACK-TBI study. *JAMA Neurol* 2021;78:982–92.
- 40 Hammond FM, Malec JF, Corrigan JD, Whiteneck GG, Hart T, Dams-O'Connor K, Novack TA, Bogner J, Dahdah MN, Eagey CB, *et al.* Patterns of functional change five to ten years after moderate-severe traumatic brain injury. *J Neurotrauma* 2021;38:1526–34.
- 41 Centers for Disease Control and Prevention, National Institutes of Health. Potential effects of a moderate or severe TBI. Available: www.cdc.gov/traumaticbraininjury/moderate-severe/potential-effects.html
- 42 Corrigan JD, Cuthbert JP, Harrison-Felix C, Whiteneck GG, Bell JM, Miller AC, Coronado VG, Pretz CR. US population estimates of health and social outcomes 5 years after rehabilitation for traumatic brain injury. *J Head Trauma Rehabil* 2014;29:E1–9.
- 43 Centers for Disease Control and Prevention, National Institutes of Health. Moderate to severe traumatic brain injury is a lifelong condition. Available: www.cdc.gov/traumaticbraininjury/pdf/Moderate_to_Severe_TBI_Lifelong-a.pdf
- 44 Pavlovic D, Pekic S, Stojanovic M, Popovic V. Traumatic brain injury: neuropathological, neurocognitive and neurobehavioral sequelae. *Pituitary* 2019;22:270–82.
- 45 Vakil E. The effect of moderate to severe traumatic brain injury (TBI) on different aspects of memory: a selective review. *J Clin Exp Neuropsychol* 2005;27:977–1021.
- 46 Azouvi P, Arnould A, Dromer E, Vallat-Azouvi C. Neuropsychology of traumatic brain injury: an expert overview. *Rev Neurol (Paris)* 2017;173:461–72.
- 47 Jourdan C, Bayen E, Pradat-Diehl P, Ghout I, Darnoux E, Azerad S, Vallat-Azouvi C, Charanton J, Aegerter P, Ruet A, *et al.* A comprehensive picture of 4-year outcome of severe brain injuries. results from the paris-TBI study. *Ann Phys Rehabil Med* 2016;59:100–6.
- 48 Belmont A, Agar N, Hugeron C, Gallais B, Azouvi P. Fatigue and traumatic brain injury. *Ann Readapt Med Phys* 2006;49:283–8.
- 49 Ponsford JL, Ziino C, Parcell DL, Shekleton JA, Roper M, Redman JR, Phipps-Nelson J, Rajaratnam SMW. Fatigue and sleep disturbance following traumatic brain injury -- their nature, causes, and potential treatments. *J Head Trauma Rehabil* 2012;27:224–33.
- 50 Sherer M, Hart T, Nick TG, Whyte J, Thompson RN, Yablon SA. Early impaired self-awareness after traumatic brain injury. *Arch Phys Med Rehabil* 2003;84:168–76.
- 51 Sherer M, Hart T, Nick TG. Measurement of impaired self-awareness after traumatic brain injury: a comparison of the patient competency rating scale and the awareness questionnaire. *Brain Inj* 2003;17:25–37.
- 52 Norup A, Mortensen EL. Prevalence and predictors of personality change after severe brain injury. *Arch Phys Med Rehabil* 2015;96:56–62.
- 53 Stéfan A, Mathé JF, group S. What are the disruptive symptoms of behavioral disorders after traumatic brain injury? A systematic review leading to recommendations for good practices. *Ann Phys Rehabil Med* 2016;59:5–17.

- 54 Schneider EB, Sur S, Raymond V, Duckworth J, Kowalski RG, Efron DT, Hui X, Selvarajah S, Hambridge HL, Stevens RD. Functional recovery after moderate/severe traumatic brain injury: a role for cognitive reserve? *Neurology* 2014;82:1636–42.
- 55 Andelic N, Bautz-Holter E, Ronning P, Olafsen K, Sigurdardottir S, Schanke AK, Sveen U, Tornas S, Sandhaug M, Roe C. Does an early onset and continuous chain of rehabilitation improve the long-term functional outcome of patients with severe traumatic brain injury? *J Neurotrauma* 2012;29:66–74.
- 56 Iwashyna TJ. Trajectories of recovery and dysfunction after acute illness, with implications for clinical trial design. *Am J Respir Crit Care Med* 2012;186:302–4.
- 57 Nagin DS. Group-based trajectory modeling: an overview. *Ann Nutr Metab* 2014;65:205–10.
- 58 Bell TM, Wang J, Nolly R, Ozdenerol E, Relyea G, Zarzaur BL. Predictors of functional limitation trajectories after injury in a nationally representative U.S. older adult population. *Ann Epidemiol* 2015;25:894–900.
- 59 Zarzaur BL, Bell T. Trajectory subtypes after injury and patient-centered outcomes. *J Surg Res* 2016;202:103–10.
- 60 Bell TM, Vetor AN, Zarzaur BL. Prevalence and treatment of depression and posttraumatic stress disorder among trauma patients with non-neurological injuries. *J Trauma Acute Care Surg* 2018;85:999–1006.
- 61 Zatzick D, Jurkovich GJ, Rivara FP, Wang J, Fan M-Y, Joesch J, Mackenzie E. A national US study of posttraumatic stress disorder, depression, and work and functional outcomes after hospitalization for traumatic injury. *Ann Surg* 2008;248:429–37.
- 62 Carr BW, Severance SE, Bell TM, Zarzaur BL. Perceived loss of social support after non-neurologic injury negatively impacts recovery. *J Trauma Acute Care Surg* 2020;88:113–20.
- 63 Prang KH, Newnam S, Berecki-Gisolf J. The impact of family and work-related social support on musculoskeletal injury outcomes: a systematic review. *J Occup Rehabil* 2015;25:207–19.
- 64 Holbrook TL, Anderson JP, Sieber WJ, Browner D, Hoyt DB. Outcome after major trauma: 12-month and 18-month follow-up results from the trauma recovery project. *J Trauma* 1999;46:765–71.
- 65 Zatzick D, Jurkovich G, Heagerty P, Russo J, Darnell D, Parker L, Roberts MK, Moodliar R, Engstrom A, Wang J, et al. Stepped collaborative care targeting posttraumatic stress disorder symptoms and comorbidity for us trauma care systems: a randomized clinical trial. *JAMA Surg* 2021;156:430–74.
- 66 deRoos-Cassini TA, Hunt JC, Geier TJ, Warren AM, Ruggiero KJ, Scott K, George J, Halling M, Jurkovich G, Fakhry SM, et al. Screening and treating hospitalized trauma survivors for posttraumatic stress disorder and depression. *J Trauma Acute Care Surg* 2019;87:440–50.
- 67 Callahan CM, Boustani MA, Unverzagt FW, Austrom MG, Damush TM, Perkins AJ, Fultz BA, Hui SL, Counsell SR, Hendrie HC. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA* 2006;295:2148–57.
- 68 Boustani MA, Sachs GA, Alder CA, Munger S, Schubert CC, Guerriero Austrom M, Hake AM, Unverzagt FW, Farlow M, Matthews BR, et al. Implementing innovative models of dementia care: the healthy aging brain center. *Aging Ment Health* 2011;15:13–22.
- 69 Khan B, Boustani M. The critical care recovery center. In: *Annual Meeting of the American Delirium Society*. 2013.
- 70 Boustani M, Unützer J, Leykum LK. Design, implement, and diffuse scalable and sustainable solutions for dementia care. *J Am Geriatr Soc* 2021;69:1755–62.
- 71 Ortiz D, Meagher AD, Lindroth H, Holler E, Gao S, Khan B, Lasiter S, Boustani M, Zarzaur B. A trauma medical home, evaluating collaborative care for the older injured patient: study protocol for a randomized controlled trial. *Trials* 2020;21:655.
- 72 Livingston DH, La Bagnara S, Sieck D, Yonclas P, Castellano C, Cho C, Walling PA, Mosenthal AC. The center for trauma survivorship: addressing the great unmet need for posttrauma center care. *J Trauma Acute Care Surg* 2020;89:940–6.