

# Respect your elders: effects of ageing on intracranial pressure monitor use in traumatic brain injury

Alexander J Schupper, Allison E Berndtson, Alan Smith, Laura Godat, Todd W Costantini

Division of Trauma, Surgical Critical Care, Burns and Acute Care Surgery, Department of Surgery, UC San Diego School of Medicine, San Diego, California, USA

## Correspondence to

Dr Todd W Costantini, Division of Trauma, Surgical Critical Care, Burns and Acute Care Surgery, Department of Surgery, UC San Diego School of Medicine, San Diego, CA 92093, USA; [tcostantini@ucsd.edu](mailto:tcostantini@ucsd.edu)

Received 19 February 2019

Revised 12 April 2019

Accepted 13 May 2019

## ABSTRACT

**Background** The Brain Trauma Foundation recommends intracranial pressure (ICP) monitor placement for patients with severe traumatic brain injury (TBI). Adherence with these guidelines in elderly patients is unknown. We hypothesized that disparities in ICP monitor placement would exist based on patient age.

**Methods** Using the National Trauma Data Bank (2010–2014), we identified patients admitted for blunt TBI with admission Glasgow Coma Scale (GCS) scores of 3–8. Patients were excluded if they had a non-Head Abbreviated Injury Scale (AIS) score  $\geq 3$ , hospital length of stay  $< 24$  hours or were discharged from the emergency department. Demographic data, ICP monitor placement, GCS, AIS-Head, Injury Severity Score, and outcome measures were collected. Propensity score matching between ICP monitor and non-ICP monitor patients was used for logistic regression and Cox multivariate regression analyses.

**Results** Of the 30 710 patients with blunt TBI with GCS scores of 3–8 included in our study, 4093 were treated with an ICP monitor. ICP monitor placement rates significantly decreased with increasing age. Multivariable analysis demonstrated that patients treated with an ICP monitor were more likely to be younger, male, have private/commercial insurance, and receive care at an institution with three or more neurosurgeons.

**Conclusion** Patients  $\geq 65$  years of age with severe blunt TBI are less likely to be treated with an ICP monitor than younger patients. Age disparities in adherence to Brain Trauma Foundation guidelines may alter the outcomes for patients with severe TBI.

**Level of evidence** Level IV.

## INTRODUCTION

Traumatic brain injury (TBI) is a considerable cause of morbidity and mortality in the USA. From 2007 to 2013, the number of TBIs sustained in the USA increased significantly.<sup>1</sup> According to the Centers for Disease Control and Prevention, TBI was responsible for approximately 2.5 million emergency department visits, 300 000 hospitalizations and 56 000 deaths in 2013 alone.<sup>1</sup> In addition to the increasing prevalence of TBI, these injuries significantly add to the national financial burden of healthcare, accounting for direct and indirect costs totaling \$60 billion in the year 2000.<sup>2</sup>

Monitoring and control of intracranial pressure (ICP) have been studied as a means of reducing secondary insults after TBI.<sup>3</sup> In the past 40 years many studies have looked at the efficacy of ICP monitor placement, but there is still a lack of

consensus on indications for use.<sup>4 5</sup> The Brain Trauma Foundation (BTF) has published guidelines for ICP monitor use, however the impact of these guidelines on patient outcomes remains uncertain.<sup>4</sup> The most recent edition of the BTF guidelines recommends the use of ICP monitoring in patients with a survivable, severe TBI (defined as Glasgow Coma Scale, or GCS, scores of 3–8) with an abnormal CT scan.<sup>6</sup> ICP monitoring may also be indicated in patients with severe TBI with a normal CT scan and two of the following features: age  $> 40$ , unilateral or bilateral motor posturing, or systolic blood pressure  $< 90$  mm Hg.<sup>6</sup> Since the publication of these guidelines, many studies have assessed their efficacy, with mixed results. A lack of consensus on the role of ICP monitoring in patients with severe TBI persists.

Elderly patients (aged 65 and older) have the highest rate of TBI-related deaths with mortality rates increasing in recent years.<sup>1</sup> This elevated mortality rate has been attributed to the higher risk of falls in this age group. Despite the high risk of mortality in elderly patients with severe TBI, the evidence for ICP monitor efficacy in this subgroup is limited.<sup>7</sup> There is a void in the literature on the effect of ICP monitoring specifically in the elderly, as well as whether or not deferring ICP monitor placement in these patients is appropriate. Our study was developed to evaluate for potential age disparities in ICP monitor placement. We hypothesized that ICP monitors would be placed less frequently in elderly patients with severe TBI compared with younger patients.

## METHODS

### Study population


To assess a widely representative population of isolated patients with TBI, we used records submitted to the National Trauma Data Bank (NTDB) from 2010 to 2014. Of the 480 347 patients in the trauma registry, we identified those patients admitted for blunt TBI with admission GCS scores between 3 and 8. Patients were subsequently excluded if they had a hospital length of stay (LOS)  $< 24$  hours, were discharged from the emergency department, transferred out to another healthcare facility, or had a non-Head Abbreviated Injury Scale (AIS) score  $\geq 3$ . Our methodology is shown in [figure 1](#).

### Covariates

Demographic data collected on our study population included patient age, sex, race and insurance

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Schupper AJ, Berndtson AE, Smith A, et al. *Trauma Surg Acute Care Open* 2019;**4**:e000306.



**Figure 1** CONSORT diagram of study methodology using the National Trauma Data Bank. AIS, Abbreviated Injury Scale; ED, emergency department; GCS, Glasgow Coma Scale; ICP, intracranial pressure; TBI, traumatic brain injury.

status. Clinical characteristics included mechanism of injury, GCS, AIS-Head, Injury Severity Score (ISS), rates of craniotomy/craniectomy and other neurosurgical procedures, types of intracranial injuries, ICP monitor placement, complications, hospital LOS, intensive care unit (ICU) LOS, and mortality. We also collected hospital teaching status, number of hospital beds and the number of neurosurgeons at the patient's hospital.

### Statistical analysis

All statistical analyses were computed via SPSS Statistics V.24, with a significance level of  $p < 0.05$ . We used logistic regression to control for demographic and clinical parameters. To directly compare the ICP monitor versus non-ICP monitor groups we used a bivariate analysis.

## RESULTS

### Demographics and clinical characteristics

The clinical characteristics of our study population can be found in [table 1](#). After applying our study inclusion and exclusion criteria, we identified 4093 patients with blunt TBI with GCS scores of 3–8 who received an ICP monitor ([table 2](#)). Patients who received an ICP monitor were significantly younger than those not receiving a monitor ( $p < 0.001$ ). Patients who received an ICP monitor also had higher ISS ( $p < 0.001$ ), higher AIS-Head

scores ( $p < 0.001$ ), and higher rates of neurosurgical procedures including craniotomy and craniectomy (both  $p < 0.001$ ). In addition, ICP monitor use was associated with a longer hospital stay (median 15 vs. 12 days,  $p < 0.001$ ), ICU stay (median 11 vs. 4 days,  $p < 0.001$ ), and more ventilator days (median 9 vs. 3 days,  $p < 0.001$ ). Patients treated with an ICP monitor were more likely to have private insurance, and a lower rate of ICP monitor placement was seen in Medicare patients ([table 3](#)). Hospitals staffed with at least three neurosurgeons favored ICP monitor placement. Finally, ICP monitor placement was associated with a higher rate of mortality (30.7% vs. 27.2%,  $p < 0.001$ , [table 2](#)).

### ICP monitor placement decreases with age

We evaluated ICP monitor use stratified by age and TBI severity. For patients with admission GCS scores of 3–8, ICP monitor placement decreased with increasing patient age ([figure 2A](#)). For example, 17% of patients aged 45–54 with admission GCS scores of 3–8 had an ICP monitor placed as compared with 10% of patients aged 65–74, and only 6% of patients aged 75–84. We also examined ICP monitor placement rates stratified by age group and AIS-Head scores ([figure 2B](#)), finding decreased rates of ICP monitor placement with increasing age for all AIS-Head groups (each  $p < 0.001$ ). The decrease in monitor placement rates by age was most significant for patients with AIS scores of 4 and 5. Next, we performed multivariable analysis to identify predictors for ICP monitor placement ([table 4](#)). Patients aged 65 and over were significantly less likely to have a monitor placed than those  $< 65$  years of age (adjusted OR 0.41, 95% CI 0.36 to 0.46). Male patients were also more likely to have an ICP monitor placed (adjusted OR 1.10, 95% CI 1.02 to 1.19). Additional predictors of ICP monitor placement included private insurance and treatment at an institution staffed by three or more neurosurgeons ([table 4](#)).

## DISCUSSION

For acute management of TBI, providers rely on literature including the BTF guidelines to support their clinical decision-making. In the 2016 fourth edition of the *Guidelines for the Management of Severe Traumatic Brain Injury*, the BTF provides a Level IIB recommendation for ICP monitoring in patients with severe TBI to reduce in-hospital and 2-week mortality.<sup>6</sup> The previous (third) edition guidelines discussed the recommendation of ICP monitoring for severe TBI (GCS scores of 3–8) with either an abnormal CT scan, or two or more of the following: ‘age over 40 years, unilateral or bilateral motor posturing, or systolic blood pressure (BP)  $< 90$  mm Hg.’<sup>8</sup> Since the third edition was published, there have not been any new studies providing further guidance on which patients should receive ICP monitoring.<sup>6</sup> In our study, we demonstrated that despite the universal inclusion of all patients with severe TBI, there is in fact an age disparity in ICP monitor placement.

Studies attempting to show a benefit after ICP monitor placement have produced mixed results. Recent studies by MacLaughlin *et al* and Agrawal *et al* have shown significant survival benefit in patients who meet BTF guideline criteria and receive an ICP monitor.<sup>9 10</sup> Unfortunately, several additional articles assessing patients who met BTF ICP monitor placement guidelines found higher mortality rates in patients who received an ICP monitor.<sup>4 11 12</sup> A recent meta-analysis by Shen *et al* of 18 studies including over 25 000 patients with severe TBI concluded that ICP monitoring significantly reduced overall mortality, hospital mortality, and 2-week and 6-month mortality rates.<sup>13</sup> However, another recent meta-analysis of patients with

**Table 1** Demographics of patients with isolated traumatic brain injury

Patients, n	30 710
Age (years, mean $\pm$ SD)	51.7 $\pm$ 20.9
Sex (male)	70.7%
ISS (IQR)	19.5 (14–26)
AIS-Head	n (%)
1	131 (0.4)
2	369 (1.2)
3	4233 (13.8)
4	12 653 (41.2)
5	13 080 (42.6)
Subarachnoid hemorrhage	14 409 (46.9)
Subdural hemorrhage	18 119 (59.0)
Epidural hematoma	2290 (7.5)
Craniotomy	4235 (13.8)
Craniectomy	1511 (4.9)
Mortality	8493 (27.7)

AIS, Abbreviated Injury Scale; ISS, Injury Severity Score.

**Table 2** Demographics and clinical characteristics (NTDB 2010–2014)

	ICP monitoring	No ICP monitoring	P value		ICP monitoring	No ICP monitoring	P value
n	4093	26 617		Alcohol use n (%)	1383 (46.6)	7575 (45.4)	0.213
Age (mean $\pm$ SD)	44.5 $\pm$ 18.4	52.8 $\pm$ 21.1	<0.001	Substance use n (%)	1175 (56.4)	6607 (54.2)	0.07
Male sex n (%)	3074 (75.1)	18 632 (70.0)	<0.001	AIS-Head n (%)			
TBI n (%)				1	2 (0.0)	129 (0.5)	0.0001
Subarachnoid hemorrhage	2189 (53.5)	12 220 (45.9)	<0.001	2	3 (0.0)	366 (1.4)	<0.0001
Subdural hemorrhage	2743 (67.0)	15 376 (57.8)	<0.001	3	225 (5.5)	4008 (15.1)	<0.0001
Epidural hematoma	419 (10.2)	1871 (7.0)	<0.001	4	1481 (36.2)	11 172 (42.0)	<0.0001
ISS (median (IQR))	22.7 (17–26)	19.0 (13–25)	<0.001	5	2373 (58.0)	10 707 (40.2)	<0.0001
Hospital days (median (IQR))	15 (8–26)	12 (6–21)	<0.001	Craniotomy n (%)	991 (24.2)	3244 (12.2)	<0.001
ICU stay (median (IQR))	11 (6–17)	4 (2–10)	<0.001	Craniectomy n (%)	582 (14.2)	929 (3.5)	<0.001
Time on ventilator (median (IQR))	9 (4–14)	3 (2–7)	<0.001	Overall mortality n (%)	1257 (30.7)	7236 (27.2)	<0.001

AIS, Abbreviated Injury Scale; ICP, intracranial pressure; ICU, intensive care unit; ISS, Injury Severity Score; NTDB, National Trauma Data Bank; TBI, traumatic brain injury.

TBI showed that ICP monitors improve prognosis, but do not affect hospital mortality rates.<sup>14</sup> This wide range of outcomes has led to the weak level of evidence regarding monitor use in recent guidelines, though the recommendation for monitor use remains. This knowledge deficit is particularly acute in trauma subpopulations, such as elderly patients, and leaves many questions unanswered. Elderly patients who suffer from a TBI have a 1-year mortality or morbidity rate of over 80%; it is unclear if this could be improved by avoiding the discrepancy in monitor use that we identify.<sup>15</sup> Recent studies focusing on specific age demographics again had mixed results, however. These include an observational study showing improved hospital and 6-month mortality with ICP monitor placement in the elderly,<sup>7</sup> as compared with a 2007–2008 NTDB study that did not find a survival benefit in patients over 55 years.<sup>5</sup>

ICP monitor placement is a safe procedure with a low-risk profile. ICP monitors are associated with some complications, including cerebrospinal fluid leak and infection, with reported rates between 0% and 5%.<sup>16–18</sup> Placement may be performed


by a wide array of specialists, with studies showing excellent outcomes with placement by trauma surgeons, neurosurgeons, general surgeons and mid-level practitioners.<sup>16–18</sup> It is unclear whether the risks of monitor placement or the risk-to-benefit ratio changes with age.

The strength of this study lies on its sampling population. Using the NTDB, we analyzed close to half a million trauma patients across the country during a 5-year period. Our sample comes from the largest national trauma registry, providing the best possible representation of trauma patients in the USA. A potential limitation to this study is the reliance on GCS as a marker of TBI. Previous studies have questioned the utility of GCS in classifying degrees of central nervous system injury, and a study by Salottolo *et al* showed that GCS can be significantly affected by age, as older patients tend to have higher GCS scores for the same severity of TBI than younger patients.<sup>19</sup> Although we used both AIS-Head and GCS as measures for brain injury severity, this potential variance does question the validity of using admission GCS scores of 3–8 as an inclusion criterion in the study as well as in the BTF guidelines. Substance abuse, in particular alcohol use, has also been shown to reduce GCS, and could potentially serve as a confounder in TBI severity.<sup>20</sup> However, rates of alcohol and substance abuse were the same in both ICP and non-ICP monitoring groups, with no statistically significant differences (table 2). To remain in accordance with current BTF guidelines, we used GCS as a primary marker of TBI severity, despite potential issues with this selection criterion. Although this study provides ample data from the hospital admission after the inciting incident, there are insufficient data

**Table 3** Insurance status and hospital data (NTDB 2010–2014)

	ICP monitoring n (%)	No ICP monitoring n (%)	P value
Insurance status			<0.001
Private/commercial	945 (23.1)	4615 (17.3)	
Self-pay	625 (15.3)	3721 (14.0)	
Blue Cross Blue Shield	258 (6.3)	1342 (5.0)	
Medicare	664 (16.2)	7962 (29.9)	
Medicaid	594 (14.5)	2996 (11.3)	
Other/unknown	384 (9.4)	2732 (10.3)	
Hospital status			0.034
Community	1467 (35.8)	9488 (35.6)	
Non-teaching	405 (9.9)	2993 (11.2)	
University	2221 (54.3)	14 136 (53.1)	
Neurosurgeons, n			<0.001
0	0 (0.0)	56 (0.2)	
1–2	285 (7.0)	2280 (8.6)	
3–5	2008 (49.1)	12 595 (47.3)	
>5	1800 (44.0)	11 686 (43.9)	

ICP, intracranial pressure; NTDB, National Trauma Data Bank.



**Figure 2** Percentage of patients who received an ICP monitor, stratified by (A) GCS and (B) AIS-Head scores. AIS, Abbreviated Injury Scale; GCS, Glasgow Coma Scale; ICP, intracranial pressure.

**Table 4** Multivariable analysis for predictors of ICP monitor placement (NTDB 2010–2014)

	OR (95% CI)	Adjusted OR (95% CI)
<b>Age (years)</b>		
<65	–	–
≥65	0.39 (0.35 to 0.42)	0.41 (0.36 to 0.46)
<b>Male sex</b>		
	1.29 (1.20 to 1.40)	1.10 (1.02 to 1.19)
<b>Insurance status</b>		
Medicaid	–	–
Private/commercial	1.03 (0.92 to 1.16)	1.14 (1.01 to 1.28)
Self-pay	0.85 (0.75 to 0.96)	0.89 (0.79 to 1.01)
Blue Cross Blue Shield	0.97 (0.83 to 1.14)	1.02 (0.86 to 1.20)
Medicare	0.42 (0.37 to 0.47)	0.72 (0.62 to 0.83)
<b>Hospital status</b>		
University	–	–
Community	0.98 (0.92 to 1.06)	1.00 (0.93 to 1.08)
Non-teaching	0.86 (0.77 to 0.96)	0.95 (0.84 to 1.06)
<b>Neurosurgeons, n</b>		
<3	–	–
≥3	1.29 (1.13 to 1.46)	1.23 (1.08 to 1.41)

ICP, intracranial pressure; NTDB, National Trauma Data Bank.

regarding long-term follow-up or functional status. This allows us to only report the immediate effects of ICP monitoring, but we are unable to comment on the long-term outcomes of this type of management.

## CONCLUSION

Patients ≥65 years of age with severe blunt TBI are less likely to be treated with an ICP monitor when compared with younger patients. Age disparities in adherence to BTF guidelines may result in worse outcomes for patients with severe TBI.

**Contributors** All authors contributed to the study design, data analysis, and article preparation.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The UC San Diego Institutional Review Board reviewed and approved this study.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open-access repository.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Taylor CA, Bell JM, Breiding MJ, Xu L. Traumatic Brain Injury-Related Emergency Department Visits, Hospitalizations, and Deaths - United States, 2007 and 2013. *MMWR Surveill Summ* 2017;66:1–16.
- Finkelstein E, Corso P, Miller T. *The incidence and economic burden of injuries in the United state*. New York: Oxford University Press, 2006.
- Becker DP, Miller JD, Ward JD, Greenberg RP, Young HF, Sakalas R. The outcome from severe head injury with early diagnosis and intensive management. *J Neurosurg* 1977;47:491–502.
- Tang A, Pandit V, Fennell V, Jones T, Joseph B, O'Keeffe T, Friese RS, Rhee P. Intracranial pressure monitor in patients with traumatic brain injury. *J Surg Res* 2015;194:565–70.
- Dang Q, Simon J, Catino J, Puente I, Habib F, Zucker L, Bukur M. More fateful than fruitful? intracranial pressure monitoring in elderly patients with traumatic brain injury is associated with worse outcomes. *J Surg Res* 2015;198:482–8.
- Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GWJ, Bell MJ, Bratton SL, Chesnut R, Harris OA, KISSOON N, *et al*. Guidelines for the management of severe traumatic brain injury, fourth edition. *Neurosurgery* 2017;80:6–15.
- You W, Feng J, Tang Q, Cao J, Wang L, Lei J, Mao Q, Gao G, Jiang J. Intraventricular intracranial pressure monitoring improves the outcome of older adults with severe traumatic brain injury: an observational, prospective study. *BMC Anesthesiol* 2015;16:35.
- Bratton SL, Chestnut RM, Ghajar J, McConnell Hammond FF, Harris OA, Hartl R, Manley GT, Nemecek A, Newell DW, Rosenthal G, *et al*. Brain trauma foundation and American Association of Neurological Surgeons. Guidelines for the management of severe traumatic brain injury. *J Neurotrauma* 2007;24:S1–S106.
- MacLaughlin BW, Plurad DS, Sheppard W, Bricker S, Bongard F, Neville A, Smith JA, Putnam B, Kim DY. The impact of intracranial pressure monitoring on mortality after severe traumatic brain injury. *Am J Surg* 2015;210:1082–7.
- Agrawal D, Raghavendran K, Schaubel DE, Mishra MC, Rajajee V. A propensity score analysis of the impact of invasive intracranial pressure monitoring on outcomes after severe traumatic brain injury. *J Neurotrauma* 2016;33:853–8.
- Shafi S, Diaz-Arrastia R, Madden C, Gentilello L. Intracranial pressure monitoring in brain-injured patients is associated with worsening of survival. *The Journal of Trauma: Injury, Infection, and Critical Care* 2008;64:335–40.
- Farahvar A, Gerber LM, Chiu Y-L, Carney N, Härtl R, Ghajar J. Increased mortality in patients with severe traumatic brain injury treated without intracranial pressure monitoring. *J Neurosurg* 2012;117:729–34.
- Shen L, Wang Z, Su Z, Qiu S, Xu J, Zhou Y, Yan A, Yin R, Lu B, Nie X, *et al*. Effects of intracranial pressure monitoring on mortality in patients with severe traumatic brain injury: a meta-analysis. *PLoS One* 2016;11:e0168901.
- Han J, Yang S, Zhang C, Zhao M, Li A. Impact of intracranial pressure monitoring on prognosis of patients with severe traumatic brain injury: a PRISMA systematic review and meta-analysis. *Medicine* 2016;95:e2827.
- Lilley EJ, Scott JW, Weissman JS, Krasnova A, Salim A, Haider AH, Cooper Z. End-of-life care in older patients after serious or severe traumatic brain injury in Low-Mortality hospitals compared with all other hospitals. *JAMA Surg* 2018;153:44–50.
- Young PJ, Bowling WM. Midlevel practitioners can safely place intracranial pressure monitors. *J Trauma Acute Care Surg* 2012;73:431–4.
- Sadaka F, Kasal J, Lakshmanan R, Palagiri A. Placement of intracranial pressure monitors by neurointensivists: case series and a systematic review. *Brain Inj* 2013;27:600–4.
- Ekeh AP, Ilyas S, Saxe JM, Whitmill M, Parikh P, Schweitzer JS, McCarthy MC. Successful placement of intracranial pressure monitors by trauma surgeons. *J Trauma Acute Care Surg* 2014;76:286–91.
- Salottolo K, Levy AS, Slone DS, Mains CW, Bar-Or D. The effect of age on Glasgow Coma Scale Score in patients with traumatic brain injury. *JAMA Surg* 2014;149:727–34.
- Rundhaug NP, Moen KG, Skandsen T, Schirmer-Mikalsen K, Lund SB, Hara S, Vik A. Moderate and severe traumatic brain injury: effect of blood alcohol concentration on Glasgow Coma Scale Score and relation to computed tomography findings. *J Neurosurg* 2015;122:211–8.