


Validity of mild TBI risk score to predict intracranial hemorrhage in cases of mild traumatic brain injury in Thailand

Kasamon Aramvanitch, Chaipayorn Yuksen , Chetsadakon Jenpanitpong, Promphet Nuanprom, Yuranan Phoonthum, Malivan Phontabtim, Sorawich Watcharakitpaisan

Emergency Medicine, Mahidol University Faculty of Medicine, Ramathibodi Hospital, Phavet, Bangkok, Thailand

Correspondence to

Dr Chaipayorn Yuksen;
Chaipool0634@hotmail.com

Received 10 February 2020

Revised 28 March 2020

Accepted 31 March 2020

ABSTRACT

Background Patients with mild traumatic brain injury (TBI) will receive a brain CT scan based on risk of injury. A previous study established a scoring system for patients with mild TBI that assigned <3 points for the low-risk group, 3–6 points for the moderate-risk group, and ≥6 points for the high-risk group. The purpose of this study was to evaluate the external validity of mild TBI risk scores for predicting intracranial hemorrhage in patients with mild TBI who had been transferred to receive a brain CT scan at the 10 nationwide CT scan-capable facilities in Thailand.

Methods The study was a retrospective cross-sectional review of patients with mild TBI who received a brain CT scan in 10 nationwide hospitals of Thailand. Risk factors were observed and points calculated for predicting mild TBI scores based on patient records. Injured patients were divided into two groups: CT scans indicating normal and abnormal brain images. After this, the accuracy of mild TBI score for predicting the presence of intracranial hemorrhage was investigated.

Results The study included a total of 999 patients, comprising 461 (46.15%) patients with abnormal brain CT scans indicating intracranial hemorrhage and 538 (53.85%) indicating no intracranial hemorrhage. In the low-risk group (mild TBI risk score <3), moderate-risk group (mild TBI risk score 3–6), and high-risk group (mild TBI risk score >6), the likelihood ratio positive of brain CT scans were 0.41, 3.53, and 77.3, respectively.

Discussion Mild TBI risk score may assist healthcare providers to select patients with mild TBI for brain CT scan referral, particularly in hospitals without CT scan facilities. In such cases, based on the proposed scoring system, immediate transfer of moderate-risk and high-risk patients with mild TBI to a CT scan-capable facility is necessary.

BACKGROUND

Annually, there are 1.4 million patients with traumatic brain injury (TBI) who visit emergency departments (EDs) in the USA. The cost of taking care of patients with TBI was US\$60 billion annually.¹ Assessment of patients with TBI in ED was affected by using the Glasgow Coma Scale (GCS) score. Mild TBI, with total GCS scores ranging from 13 to 15, the largest number of patients with TBI, accounted for 80%.² Only 15% of patients with a GCS score of 15 demonstrated an abnormality

on CT and only 1% of these patients required cranial surgery.^{3,4} This group of patients was usually neglected from observation and closed monitoring when comparing with patients with moderate and severe TBI.⁵ A GCS score of 13 was associated with 100% risk for intracranial hemorrhage.⁶

Patients with mild TBI were divided into three groups: a high-risk group, a moderate-risk group, and a low-risk group.^{3,7–9} Clinical factors for high-risk group included male gender, age ≥60 years, baseline GCS score, clinical signs of skull fracture, clinical signs of basilar skull fracture, a GCS score drop ≥2 points, post-traumatic vomiting, focal neurological signs, seizure, headache, severe headache (visual analog scale score >7), post-traumatic amnesia (<1 hour), transient loss of consciousness, large extracranial hematoma/severe maxillofacial injury, coagulopathy, trauma mechanisms, and drug/alcohol intoxication.¹⁰ Per the guidelines developed by the Royal College of Surgeons of Thailand and the College of Neurological Surgeons of Thailand, patients with mild TBI who are considered high risk should undergo a brain CT scan within 1 hour of ED admission.

The study in Ramathibodi Hospital, a university-affiliated super tertiary care hospital in Bangkok, Thailand, was conducted to develop the clinical prediction score to predict intracranial hemorrhage in patients with mild TBI. The mild TBI risk scores were categorized into three groups: scores <3 (low risk), scores 3–6 (moderate risk), and scores >6 (high risk). The likelihood ratio of positive head CT scan was 20.61 (95% CI 12.74 to 33.33) in the high-risk group.¹⁰

Therefore, the objective of this study was to evaluate the external validity of mild TBI risk scores for predicting intracranial hemorrhage in patients with mild TBI who had been transferred to receive a brain CT scan at the 10 nationwide CT scan-capable facilities in Thailand.

PATIENTS AND METHODS

The study was a retrospective cross-sectional review of patients with isolated mild TBI who had been transferred to receive a brain CT at the 10 nationwide CT scan-capable facilities in Thailand. The inclusion criteria were age ≥15 years. Those patients who had been transferred to receive a brain CT scan without medical records were

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Aramvanitch K, Yuksen C, Jenpanitpong C, et al. *Trauma Surg Acute Care Open* 2020;**5**:e000453.

excluded. The study period was between June and December 2016.

The data of eligible patients were recorded, including baseline characteristic and potential risk factors for intracranial hemorrhage. Risk factors included post-traumatic vomiting more than two times (1.5 points), severe headache (7.5 points), transient loss of consciousness (3 points), post-traumatic amnesia (1 point), focal neurological signs (4 points), clinical signs of skull fracture (6 points), and base of skull fracture (8 points).¹⁰ All eligible patients were divided into two groups according to CT scan results: CT positive, indicating intracranial hemorrhage; and CT negative, indicating no intracranial hemorrhage. The outcome of this study was accuracy of mild TBI risk score for predicting the presence of intracranial hemorrhage.

Statistical analysis

Baseline characteristics of eligible patients were compared for difference between two groups divided by brain CT results. Exact probability test was used to analyze the difference in categorical variables, whereas an independent t-test or Wilcoxon rank-sum test was used in continuous variables.

Assessing discrimination of studied risk factors for positive CT results presented as an area under receiving operating characteristic (AuROC) curve, 95% CI, and significance level (p value). The patient's risk of intracranial hemorrhage was categorized into low, moderate, and high risk according to mild TBI risk score calculation. Diagnostic indices including sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and likelihood ratio positive (LR+) were examined to assess the validity of mild TBI score.

Results

In this study, there were 999 patients with mild TBI who had been transferred to 10 nationwide CT scan-capable facilities to receive a brain CT scan. Of those, 461 patients (46%) indicated the presence of intracranial hemorrhage (CT positive), and 538 patients (54%) resulted in normal brain imaging (CT negative). According to the type of incidents, motorcycle crash and assault had higher result significantly in CT positive (237 (51.41%) vs 137 (25.46%), $p < 0.001$ and 38 (8.24%) vs 16 (2.97%), $p < 0.001$, respectively) (table 1).

Regarding the univariable analysis, there were nine potential risk factors that had significantly higher proportion in positive CT findings and high discriminative performance (AuROC) (table 2). The potential risk factors are suspicion of skull fracture (20.82% vs 0.37%, $p < 0.001$) (AuROC curve of 0.60 and 95% CI 0.57 to 0.63), neurological deficit (12.36% vs 5.39%, $p < 0.001$) (AuROC curve of 0.60 and 95% CI 0.53 to 0.57), experiencing of post-traumatic amnesia (58.83% vs 24.35%, $p < 0.001$) (AuROC curve of 0.66 and 95% CI 0.63 to 0.69), and history of loss of consciousness (67.68% vs 16.17%, $p < 0.001$) (AuROC curve of 0.76 and 95% CI 0.73 to 0.78).

The accuracy of mild TBI risk score for predicting intracranial hemorrhage was investigated (table 3). In low-risk patients, sensitivity was 8.24%, specificity 79.74%, PPV 25.85%, NPV 50.35%, and LR+ 0.41. In moderate-risk patients, sensitivity was 57.05%, specificity 83.83%, PPV 75.14%, NPV 69.50%, and LR+ 3.53. For those of high risk, sensitivity was 28.63%, specificity 99.63%, PPV 98.51%, NPV 61.97%, and LR+ 77.3.

The discrimination performance of mild TBI risk score for predicting intracranial hemorrhage was presented as AuROC curve of 0.89 and 95% CI 0.87 to 0.91.

Table 1 Baseline prognostic factors of patients with mild TBI categorized by brain CT scan results

| Characteristics | CT positive (n=461) | | CT negative (n=538) | | P value |
|---|------------------------|-------|------------------------|-------|---------|
| Age (mean±SD) | 46.3±24.1 | | 54.9±25.0 | | <0.001 |
| Admission systolic blood pressure (mean±SD) | 125.3±34.8 | | 110.1±28.8 | | 0.225 |
| ED triage level | | | | | |
| Level 1 | 14 | 3.1% | 9 | 1.7% | 0.416 |
| Level 2 | 247 | 54.3% | 211 | 39.5% | |
| Level 3 | 186 | 40.9% | 296 | 55.4% | |
| Level 4 | 7 | 1.5% | 17 | 3.2% | |
| Level 5 | 1 | 0.2% | 1 | 0.2% | |
| Type of injury | | | | | |
| Motorcycle crash | 237 | 51.4% | 137 | 25.5% | <0.001 |
| Motor vehicle crash | 31 | 6.7% | 23 | 4.3% | 0.094 |
| Falling | 108 | 23.4% | 286 | 53.2% | <0.001 |
| Blunt object | 24 | 5.2% | 26 | 4.8% | 0.884 |
| Assault | 38 | 8.2% | 16 | 3.0% | <0.001 |
| Fainting/syncope | 28 | 6.1% | 43 | 8.0% | 0.267 |
| Laboratory results* | | | | | |
| Platelet (×10 ³) | 222.6 | 836.8 | 243.2 | 119.2 | 0.088 |
| INR | 1.1 | 0.3 | 1.1 | 0.1 | 0.785 |
| Type of TBI | | | | | |
| EDH | 105 | 22.8% | | | |
| SDH | 91 | 19.7% | | | |
| SAH | 88 | 19.1% | | | |
| Brain contusion | 72 | 15.6% | | | |
| IPH | 105 | 23.0% | | | |
| Hospital stay (days) | 4.5±6.9 | | 2.7±3.1 | | <0.001 |
| Disposition | | | | | |
| Discharge | 446 | 96.8% | 536 | 99.6% | <0.001 |
| Dead | 15 | 3.3% | 2 | 0.4% | |

*Laboratory results: platelet concentration and INR reported as mean and SD. ED, emergency department; EDH, epidural hematoma; INR, international normalized ratio; IPH, intraparenchymal hemorrhage; SAH, subarachnoid hemorrhage; SDH, subdural hematoma; TBI, traumatic brain injury.

Discussion

According to the guidelines developed by the Royal College of Surgeons of Thailand and the College of Neurological Surgeons of Thailand, patients with mild TBI can be categorized into three groups: high-risk, moderate-risk, and low-risk groups. Of those, patients with mild TBI who are considered high risk should undergo an expedited brain CT.³⁻⁷⁻⁹

Delays in obtaining this CT scan in high-risk patients with mild TBI may lead to worse outcomes including increased disability and mortality.¹⁰ In contrast, patients with mild TBI who are low risk are unlikely to have an intracranial hemorrhage and should avoid a CT scan to prevent unnecessary radiation and costs.¹¹

Regarding the study, overall, 46% of eligible patients who underwent brain CT demonstrated intracranial hemorrhage. For low-risk patients with mild TBI, the LR+ of positive CT findings (intracranial hemorrhage) was only 0.41. Management of these patients for non-CT scan-capable facilities should be neurological observation and transport to appropriate facility if neurological deficit develops.

Patient with moderate risk were found that the LR+ increased to 3.35 times and increased to 77.3 times for patients with high risk. Management of these patients for non-CT scan-capable facilities should be rapid transport to the closest appropriate

Table 2 Potential risk factors of patients with mild TBI categorized by brain CT scan results

| Risk factors | CT positive (n=461) | | CT negative (n=538) | | P value | AuROC (95% CI) |
|--------------------------------|---------------------|------|---------------------|------|---------|---------------------|
| | N | % | N | % | | |
| Gender—male | 176 | 38.2 | 226 | 42.0 | 0.220 | 0.52 (0.49 to 0.55) |
| Age >60 years | 145 | 31.5 | 306 | 56.9 | <0.001 | 0.37 (0.34 to 0.40) |
| Admission GCS score | | | | | | |
| GCS 15 | 385 | 83.5 | 490 | 91.1 | 0.001 | 0.46 (0.43 to 0.49) |
| GCS 14 | 51 | 11.1 | 33 | 6.1 | | |
| GCS 13 | 25 | 5.4 | 15 | 2.8 | | |
| Skull fracture | 96 | 20.8 | 2 | 0.4 | <0.001 | 0.60 (0.57 to 0.63) |
| Basilar skull fracture | 10 | 2.2 | 0 | 0 | <0.001 | 0.51 (0.48 to 0.54) |
| GCS score drop ≥ 2 points | 20 | 4.3 | 8 | 1.5 | 0.007 | 0.51 (0.48 to 0.55) |
| Penetrating skull injury | 3 | 0.7 | 0 | 0 | 0.098 | 0.50 (0.47 to 0.53) |
| Vomiting ≥ 2 times | 40 | 8.7 | 47 | 8.7 | 0.999 | 0.50 (0.47 to 0.53) |
| Neurological deficit | 57 | 12.4 | 29 | 5.4 | <0.001 | 0.53 (0.50 to 0.57) |
| Post-traumatic seizure | 17 | 3.7 | 14 | 2.6 | 0.363 | 0.51 (0.47 to 0.54) |
| Headache | 164 | 35.6 | 131 | 24.4 | <0.001 | 0.56 (0.53 to 0.59) |
| Severe headache | 33 | 7.2 | 0 | 0 | <0.001 | 0.53 (0.50 to 0.57) |
| Post-traumatic amnesia | 262 | 58.8 | 131 | 24.4 | <0.001 | 0.66 (0.63 to 0.69) |
| History of LOCs | 312 | 67.7 | 87 | 16.2 | <0.001 | 0.76 (0.73 to 0.78) |
| Scalp/facial hematoma | 85 | 18.4 | 41 | 7.6 | <0.001 | 0.55 (0.52 to 0.59) |
| Intoxication | 89 | 25.9 | 35 | 22.6 | 0.502 | 0.52 (0.47 to 0.56) |

AuROC, area under receiving operating characteristic curve; GCS, Glasgow Coma Scale; LOC, loss of consciousness; TBI, traumatic brain injury.

Table 3 Diagnostic indices of mild TBI risk score categorized by risk category

| Risk factors | CT positive | CT negative | Sensitivity (%) | Specificity (%) | PPV (%) | NPV (%) | LR+ |
|---------------|-------------|-------------|-----------------|-----------------|---------|---------|------|
| Low risk | 38 | 109 | 8.2 | 79.7 | 25.9 | 50.4 | 0.4 |
| Moderate risk | 263 | 87 | 57.1 | 83.8 | 75.1 | 69.5 | 3.5 |
| High risk | 132 | 2 | 28.6 | 99.6 | 98.5 | 62.0 | 77.3 |

Low risk, mild TBI score <3 points; moderate risk, mild TBI score 3–6 points; high risk, mild TBI score ≥ 7 points.

LR+, likelihood ratio positive; 0.89 and 95% CI 0.87 to 0.91; NPV, negative predictive value; PPV, positive predictive value; TBI, traumatic brain injury.

facility to provided them a brain CT scan. And for CT scan-capable facilities, patients with mild TBI should be prioritized for receiving brain imaging based on their risk category.

The study had a number of limitations including the retrospective nature of the data reviewed. Some of the background data were excluded because they were incomplete. Not all high-risk patients with mild TBI were transferred for a CT scan.

In conclusion, for patients with mild TBI, a risk score can help predict the presence of intracranial hemorrhage which can be used to determine and prioritize the patients who require a brain CT. Those patients with mild TBI with high-risk and moderate-risk scores should undergo immediate CT scans.

Collaborators No.

Contributors KA and CY conceptualised the research idea, analyzed and interpreted the data. CJ participated in data collection and article writing. PN, YP, SW, and MP participated in data collection.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved by the institutional review board, Faculty of Medicine, Ramathibodi Hospital Mahidol University, Thailand (MURA2017/532).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. Deidentified participant data.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Chaiyaporn Yuksen <http://orcid.org/0000-0002-4890-7176>

REFERENCES

- 1 Faul MXL, Wal MM, Coronado VG. US Department of Health and Human Services. *Traumatic brain injury in the United States: emergency department visits, hospitalizations, and deaths*. Atlanta, GA: CDC, 2010:2002–6.
- 2 Levin HS. Outcome from mild head injury. In: Narayan RK, Wilberger JE, Povlishock JT, eds. *Neurotrauma*. New York: McGraw-Hill, 1996:755–65.
- 3 Haydel MJ, Preston CA, Mills TJ, Luber S, Blaudeau E, DeBlieux PMC, et al. Indications for computed tomography in patients with minor head injury. *N Engl J Med Overseas Ed* 2000;343:100–5.
- 4 Stiell IG, Clement CM, Rowe BH, Schull MJ, Brison R, Cass D, Eisenhauer MA, McKnight RD, Bandiera G, Holroyd B, et al. Comparison of the Canadian CT head rule and the New Orleans criteria in patients with minor head injury. *JAMA* 2005;294:1511–8.
- 5 Galbraith S. Misdiagnosis and delayed diagnosis in traumatic intracranial haematoma. *Br Med J* 1976;1:1438–9.
- 6 Yuksen C, Sittichanbuncha Y, Patumanond J, Muengtawepong S, Aramvanich K, Supamas A, Sawanyawisuth K. Clinical factors predictive for intracranial hemorrhage in mild head injury. *Neurol Res Int* 2017;2017:5385613–5.

- 7 Smits M, Dippel DWJ, Steyerberg EW, de Haan GG, Dekker HM, Vos PE, Kool DR, Nederkoorn PJ, Hofman PAM, Twijnstra A, *et al.* Predicting intracranial traumatic findings on computed tomography in patients with minor head injury: the CHIP prediction rule. *Ann Intern Med* 2007;146:397–405.
- 8 Kuppermann N, Holmes JF, Dayan PS, Hoyle JD, Atabaki SM, Holubkov R, Nadel FM, Monroe D, Stanley RM, Borgialli DA, *et al.* Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *Lancet* 2009;374:1160–70.
- 9 Brenner DJ. Estimating cancer risks from pediatric CT: going from the qualitative to the quantitative. *Pediatr Radiol* 2002;32:228–31.
- 10 Yuksen C, Sittichanbuncha Y, Patumanond J, Muengtaweepongs S, Sawanyawisuth K. Clinical predictive score of intracranial hemorrhage in mild traumatic brain injury. *Ther Clin Risk Manag* 2018;14:eCollection:213–8.
- 11 Smits M, Dippel DWJ, de Haan GG, Dekker HM, Vos PE, Kool DR, Nederkoorn PJ, Hofman PAM, Twijnstra A, Tanghe HLJ, *et al.* Minor head injury: guidelines for the use of CT—a multicenter validation study. *Radiology* 2007;245:831–8.