

Neurological outcomes after traumatic cardiopulmonary arrest: a systematic review

Daniel Shi,¹ Christie McLaren,¹ Chris Evans²

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tsaco-2021-000817>).

¹School of Medicine, Queen's University, Kingston, Ontario, Canada

²Emergency Medicine, Queen's University, Kingston, Ontario, Canada

Correspondence to

Daniel Shi; dshi@qmed.ca

This study was presented as an oral presentation at the Annual Meeting of the Trauma Association of Canada in a virtual format on April 12, 2021.

Received 3 August 2021

Accepted 10 October 2021

ABSTRACT

Background Despite appropriate care, most patients do not survive traumatic cardiac arrest, and many survivors suffer from permanent neurological disability. The prevalence of non-dismal neurological outcomes remains unclear.

Objectives The aim of the current review is to summarize and assess the quality of reporting of the neurological outcomes in traumatic cardiac arrest survivors.

Data sources A systematic review of Embase, Medline, PubMed, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and ProQuest databases was performed from inception of the database to July 2020.

Study eligibility criteria Observational cohort studies that reported neurological outcomes of patients surviving traumatic cardiac arrest were included.

Participants and interventions Patients who were resuscitated following traumatic cardiac arrest.

Study appraisal and synthesis methods The quality of the included studies was assessed using ROBINS-I (Risk of Bias in Non-Randomized Studies - of Interventions) for observational studies.

Results From 4295 retrieved studies, 40 were included (n=23 644 patients). The survival rate was 9.2% (n=2168 patients). Neurological status was primarily assessed at discharge. Overall, 45.8% of the survivors had good or moderate neurological recovery, 29.0% had severe neurological disability or suffered a vegetative state, and 25.2% had missing neurological outcomes. Seventeen studies qualitatively described neurological outcomes based on patient disposition and 23 studies used standardized outcome scales. 28 studies had a serious risk of bias and 12 had moderate risk of bias.

Limitations The existing literature is characterized by inadequate outcome reporting and a high risk of bias, which limit our ability to prognosticate in this patient population.

Conclusions or implications of key findings Good and moderate neurological recoveries are frequently reported in patients who survive traumatic cardiac arrest. Prospective studies focused on quality of survivorship in traumatic arrest are urgently needed.

Level of evidence Systematic review, level IV.

PROSPERO registration number CRD42020198482.

BACKGROUND

Traumatic cardiac arrest (TCA) occurs due to severe injury, most commonly from traumatic brain injury and hemorrhage.¹ Despite appropriate care, TCA has been associated with extremely low survival rates, with 2% of patients surviving to hospital discharge.² Some authors have even concluded

that resuscitation of patients with TCA is futile and costly.³ However, recent data from prospectively registered trauma systems in England, Spain, and North America have suggested that outcomes from TCA may be better than previously expected, with overall survival rates between 5.7% and 7.5%.⁴⁻⁶ Advances in damage control resuscitation and our understanding of its pathophysiology have led to improvements in the contemporary management of TCA, which is at least partially responsible for the observed increase in overall survival.⁷

The prognostication of patients with TCA is an important consideration for patients, families, and health providers that initiate resuscitative efforts, as after survival the main treatment goal is a favorable neurological outcome.^{3,8} Poor outcomes after cardiac arrest of any etiology have been attributed to hypoxic-ischemic brain injury.⁹ The extent of this brain injury is an important predictor of unfavorable neurological outcomes, which are defined by death from neurological cause, persistent vegetative state, or severe neurological disability.⁹ Most studies examining cardiac arrest outcomes use the Cerebral Performance Categories (CPCs) or the Glasgow Outcome Scale (GOS) to report neurological status.⁹ Other scales that have demonstrated value in assessing the neurological outcome of survivors include the modified Rankin Scale (mRS), the Extended Glasgow Coma Scale (GOSE), and the Functional Independence Measure (FIM).¹⁰⁻¹² Despite the existence of several validated measures, in many neuroprognostication studies, neurological outcomes are generally dichotomized as “good” or “poor”, with no consensus on how a poor outcome is defined.⁹ The neurological outcome of these patients remains unclear in the current literature. Among TCA survivors, residual neurological deficits have generally been found to be severe and disabling in small observational studies.^{13,14} However, a systematic review in 2012 found that more than half of TCA survivors either make a full neurological recovery or have moderate deficits.¹⁵

Although previous literature has focused on summarizing the proportion of survivors and identifying resuscitation techniques to reduce mortality in traumatic arrest, minimal evidence exists that assesses the quality of neurological outcome data. Hence, we performed a systematic review to summarize the neurological outcomes of patients who survive TCA. Particular attention was placed on the variation of reporting and definitions of these outcomes between studies.

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Shi D, McLaren C, Evans C. *Trauma Surg Acute Care Open* 2021;**6**:e000817.

METHODS

This review was conducted in accordance with the *Cochrane Handbook for Systematic Reviews of Interventions* and followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement.¹⁶ The protocol for this study was registered with the International Prospective Register of Systematic Reviews.

Inclusion criteria

Eligible studies included observational cohort studies that enrolled patients who experienced cardiopulmonary arrest following trauma and reported neurological outcomes. Studies examining pediatric patients were included. We excluded case reports, case series studies, reviews, and animal studies. Studies published in the English language were included. There were no restrictions on the length of follow-up, geographical location, or publication date.

Study selection and data abstraction

A systematic search of Embase, Medline, PubMed, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and ProQuest was conducted from the inception of the database to July 2020. A librarian with methodological and content expertise was consulted to create the search strategies (online supplemental table 1). Medical subject headings and Emtree headings were used in their respective databases.

Two reviewers (DS and CM) independently screened the title and abstract of all included studies. Duplicate studies were

removed using Covidence.¹⁷ Studies that met the inclusion criteria were reviewed in full text by the same two reviewers independently (figure 1). Disagreements were resolved by consensus or by the decision of a third independent reviewer (CE). Interobserver agreements for the title/abstract and full text stages were calculated using Cohen's κ statistics.¹⁸

Data were abstracted by two independent reviewers using a piloted data abstraction form (Microsoft Excel). The primary outcome was the neurological status of TCA survivors which was abstracted from each study. Additionally, the methods used to assess neurological outcome, the mechanism of trauma, and the time point when neurological status was measured were abstracted.

Data analysis

Patients from the included studies were placed into one of four categories to describe their neurological outcome: good, moderate, poor, or vegetative. A "good" neurological outcome was defined as a full neurological recovery or having minor deficits. A "moderate" outcome was defined as having neurological disability that partially affects daily activities but having full independence. A "poor" outcome was defined as requiring any level of dependency or personal assistance in daily living. A "vegetative" outcome was defined by unresponsiveness to external stimuli and a decreased level of awareness.

The proportion of patients with "favorable" neurological outcomes (ie, patients with good or moderate outcome) versus




Figure 1 PRISMA diagram. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

patients with “unfavorable” neurological outcomes (ie, patients with poor or vegetative outcome) was also calculated.

Quality assessment

The quality of included studies was assessed using ROBINS-I (Risk of Bias in Non-Randomized Studies - of Interventions) tool for observational studies. The overall risk of bias was rated as critical, serious, moderate, or low. Quality assessment was completed by two independent reviewers (DS and CM), with all conflicts resolved by consensus or discussion with a third reviewer (CE).

RESULTS

Study characteristics

After the removal of duplicates, the literature search generated 4295 citations for title and abstract review. One hundred and eighty-one qualified for full-text screening, of which 40 were included in our analysis (figure 1). The title and abstract screening showed almost perfect agreement ($\kappa=0.83$), and similar agreement was found for the full-text screening ($\kappa=0.87$). Of the included studies, 35 were retrospective cohort studies and 5 were prospective cohort studies (table 1). The included studies were published between 1983 and 2019, with 12 studies published in the past 5 years.

Table 1 Characteristics of included studies

Study identification	Design	Age group	Sample size	Survivors (n)	Mortality (%)	Blunt trauma	Penetrating trauma	Other/unknown trauma type
Alanezi ⁴⁵ 2004	Prospective cohort	Mixed	50	2	96.0	4	46	0
Barnard ⁴ 2017	Retrospective cohort	Mixed	705	53	92.5	601	104	0
Beck ⁴⁶ 2016	Retrospective cohort	Mixed	1354	9	99.3	869	226	259
Calkins ³⁴ 2002	Retrospective cohort	Pediatric	25	2	92.0	25	0	0
Capizzani ³⁵ 2010	Retrospective cohort	Pediatric	30	6	80.0	21	0	9
Chia ⁴⁷ 2017	Retrospective cohort	Mixed	1554	38	97.6	NR	NR	1554
Chiang <i>et al</i> ⁴⁸ 2015	Retrospective cohort	Adult	514	20	96.1	388	0	126
Chien ⁴⁹ 2016	Retrospective cohort	Adult	396	9	97.7	197	11	188
David ⁵⁰ 2007	Retrospective cohort	Adult	268	6	97.8	NR	NR	268
Deasy ⁵¹ 2012	Retrospective cohort	Mixed	2187	28	98.7	NR	NR	2187
Di Bartolomeo ¹⁹ 2005	Retrospective cohort	Adult	181	2	89.0	181	0	0
Djarv ⁵² 2018	Retrospective cohort	Mixed	1774	65	96.3	NR	NR	1774
Duchateau ⁵³ 2017	Retrospective cohort	Adult	88	10	88.6	77	11	0
Evans ⁵ 2016	Retrospective cohort	Mixed	2300	145	92.7	1547	736	270
Falcone ⁵⁴ 1995	Retrospective cohort	Adult	320	6	98.1	285	36	0
Fisher ⁵⁵ 1999	Retrospective cohort	Pediatric	65	1	98.5	65	0	0
Graesner ⁵⁶ 2011	Retrospective cohort	Mixed	814	221	72.9	597	52	165
Hillman ⁵⁷ 2016	Retrospective cohort	Pediatric	27	4	85.2	4	7	16
Huber-Wagner ⁵⁸ 2007	Retrospective cohort	Mixed	757	130	82.8	714	43	0
Keller ³⁷ 2013	Retrospective cohort	Mixed	448	37	91.7	NR	NR	NR
Kleber ²³ 2014	Prospective cohort	Mixed	71	15	79.9	40	12	19
Lawhon ²⁰ 1995	Retrospective cohort	Mixed	47	2	95.7	NR	NR	47
Lin ⁵⁹ 2016	Retrospective cohort	Pediatric	388	38	91.2	365	23	0
Love ⁶⁰ 2016	Retrospective cohort	Mixed	237	7	97.0	165	72	0
Lundy ⁶¹ 2011	Retrospective cohort	Mixed	309	125	59.5	NR	NR	309
Molina ⁶² 2008	Retrospective cohort	Mixed	94	8	91.5	0	94	0
Mollberg ⁶³ 2011	Retrospective cohort	Adult	294	1	99.7	90	204	0
Moore ⁴⁴ 2011	Prospective cohort	Mixed	NR	56	NR	5	51	NR
Moore ⁶⁴ 2016	Prospective cohort	Adult	1708	106	93.8	820	888	0
Murphy ⁶⁵ 2010	Retrospective cohort	Pediatric	169	28	83.4	151	7	11
Perron ⁶⁶ 2001	Retrospective cohort	Pediatric	729	184	74.8	505	81	143
Pickens ⁶⁷ 2005	Retrospective cohort	Mixed	184	14	92.4	90	94	0
Powell ⁶⁸ 2004	Retrospective cohort	Mixed	959	62	93.5	11	51	897
Rabinovici ⁶⁹ 2014	Retrospective cohort	Mixed	67	9	86.6	17	50	0
Shimazu ⁷⁰ 1983	Retrospective cohort	Mixed	267	7	97.4	217	50	0
Stratton ⁷¹ 1998	Retrospective cohort	Mixed	879	9	99.9	382	497	0
Tarmey ⁴³ 2011	Prospective cohort	Adult	52	4	92.3	0	17	35
Vassallo ⁷² 2019	Retrospective cohort	Pediatric	129	7	94.6	110	19	0
Zwingmann ⁷³ 2015	Retrospective cohort	Pediatric	152	43	71.7	145	7	0
Zwingmann ²¹ 2016	Retrospective cohort	Mixed	3052	649	78.7	NR	NR	3052

NR, not reported.

Patient characteristics

There were 23 644 patients included, with 2168 (9.2%) surviving to hospital discharge. The mortality rate was 90.8%. Blunt trauma (n=8687 patients) and penetrating trauma (n=3489 patients) were the main mechanisms of traumatic arrest, whereas the remaining trauma mechanisms were not classified (n=11 468) (table 1). Nine studies examined pediatric patients only (n=1714 patients), 9 studies examined adults only (n=3821 patients), and 22 studies examined a mixed population (n=18 109 patients) without providing information of how many pediatric patients were included.

Neurological outcome reporting

Seventeen studies qualitatively described their neurological outcome based on patient disposition after discharge (eg, home, nursing care, or rehabilitation) and the level of assistance required in daily activities. Thirteen studies used the GOS and reported the number of patients in each GOS category (GOS 1–5). Eight studies used the CPC and one study used the Pediatric Cerebral Performance Categories (PCPC) to assess neurological outcome.

Seven studies dichotomized neurological outcomes into “good” or “poor” categories using the GOS, CPC, or PCPC scales. Six of the seven studies considered a “good” neurological outcome as patients with CPC 1 and 2 or GOS 4 and 5 (ie, good, mild, or moderate deficits) and a “poor” neurological outcome as those with CPC 3 and 5 or GOS 1 and 3 (ie, severe or vegetative). One study used the PCPC and defined a “good” outcome as PCPC 1 and 2 (good or mild deficits) and a “poor” outcome as PCPC 3 and 5 (moderate or severe deficits, or vegetative).

One study used the FIM to assess neurological status. This study reported an average FIM score for the survivors and the number of patients who had obtained the lowest score (required assistance in daily life). The functional status of the other patients was not reported.

Neurological status was recorded primarily at patient discharge, but some studies reported outcomes up to a 4-year follow-up period.

Overall neurological outcomes

The neurological outcomes of individual studies are reported in table 2. Based on the 32 studies (n=1507 patients) that reported the number of patients for each neurological outcome category, 538 (35.7%) had good outcomes, 392 (26.0%) had moderate outcomes, 408 (27.1%) had poor outcomes, and 169 (11.2%) were vegetative.

In six studies (n=115 patients) that reported dichotomous outcomes, 62 patients (53.9%) had “good” outcomes and 53 patients (46.1%) had “poor” outcomes. In the one study (n=38 patients) that used the PCPC, 12 (31.6%) had “good” outcomes and 26 (68.4%) had “poor” outcomes. One study (n=184 patients) that used the FIM reported an average score of 38.9 (range: 18–126), 7 patients with severe deficits and 177 patients that had either good recovery or moderate or severe deficits.

Based on all 40 studies (n=2168 patients) included in this review, 992 (45.8%) had favorable outcomes, 630 (29.0%) had unfavorable outcomes, and 546 (25.2%) were missing. The results of the studies that used the PCPC and the FIM were included in the missing category, as the neurological status of the survivors was unclear and could not be categorized.

Neurological outcomes for pediatric studies

Of the seven studies (n=87 patients) that examined only pediatric populations and reported the number of patients for each

neurological outcome category, 37 (42.5%) had good outcomes, 19 (21.8%) had moderate outcomes, 18 (20.7%) had poor outcomes, and 13 (14.9%) were vegetative. Hence, 56 patients (64.4%) had favorable outcomes and 31 (35.6%) had unfavorable outcomes.

Neurological outcomes for adult studies

Of the nine studies (n=164 patients) that examined only adult populations, 119 patients (72.6%) had favorable outcomes, 41 (25.0%) had unfavorable outcomes, and 4 (2.4%) had missing outcomes. Of the six studies (n=129 patients) that reported the number of patients for each neurological outcome category, 87 (67.4%) had good outcomes, 15 (11.6%) had moderate outcomes, 6 (4.7%) had poor outcomes, 17 (13.2%) were vegetative, and 4 (3.1%) had missing outcomes.

Assessment of quality

The quality of all included studies (n=40 studies) was assessed. Twenty-eight studies were at a serious risk of bias and 12 studies were at a moderate risk of bias (table 3). In general, the studies were well reported. The confounding domain was the primary source of bias, as most studies did not consider many potential confounders, including age, resuscitation technique, and type of trauma. The outcome measurement domain was also a significant source of bias, as most studies used physician-reported neurological outcomes, which were described qualitatively in the studies. Hence, these studies were rated as a serious risk of bias in this domain. Finally, there was moderate to serious risk of bias for selective reporting, as some studies did not report outcomes for each neurological category.

DISCUSSION

The current review is the first study to focus on neurological outcomes across a large patient population (n=23 644 patients). We report a survival rate of 9.2%, which is one of the most optimistic findings to date for outcomes following TCA. In the current review, favorable neurological outcomes were frequently reported (45.8% with full recovery or moderate disability), suggesting that outcomes from TCA may be more favorable than previously expected.³

Previous literature suggests that prognosis after traumatic arrest is extremely poor. Many studies reported only a small number of survivors, all with severe neurological disability.^{3 14 19 20} However, in 2012, new studies were conducted and a systematic review found that good and moderate neurological outcomes were reported in 57.4% of survivors.¹⁵ A large retrospective study published in 2016 supports these findings and reported good and moderate neurological outcomes in up to 75.0% of survivors.²¹ Despite these findings, neurological outcomes continue to be debated, as newer studies with larger numbers of survivors report good and moderate outcomes in only 4.3% to 27.0% of survivors.^{22 23} The findings of the current review suggest that those who survive traumatic arrest may have a favorable prognosis. The observed improvement in neurological outcomes is likely connected to novel advances in damage control resuscitation and refinement of treatment of guidelines.⁷

A secondary aim of this review was to assess the quality of the reporting of neurological outcomes. In most of the included studies, neurological status was qualitatively described. As the exact deficits were often not described, it was difficult to quantify the extent of the impairments and classify the neurological outcomes of these patients. Alternatively, patient disposition (eg, home, nursing care, rehabilitation) and dependence

Table 2 Neurological outcomes of included studies

Study identification	Neurological scale	Time point	Qualitative outcomes				
			Good	Moderate	Poor	Vegetative	Missing
Alanezi ⁴⁵ 2004	Qualitative	Discharge	1	0	1	0	0
Barnard ⁴ 2017	GOS	Discharge	24*	N/A	9†	N/A	20
Beck ⁴⁶ 2016	CPC	Discharge	6*	N/A	3	0	0
Calkins ³⁴ 2002	Qualitative	Discharge	2	0	0	0	0
Capizzani ³⁵ 2010	Qualitative	Discharge	2	1	3	0	0
Chia ⁴⁷ 2017	CPC	Discharge	15*	N/A	23†	N/A	0
Chiang ⁴⁸ 2015	CPC	Discharge	12*	N/A	8†	N/A	0
Chien ⁴⁹ 2016	CPC	Discharge	3*	N/A	6†	N/A	0
David ⁵⁰ 2007	CPC	Discharge	2*	N/A	4†	N/A	0
Deasy ⁵¹ 2012	GOS	1 year	0	2	2	1	23
Di Bartolomeo ¹⁹ 2005	GOS	4 years	0	0	2	0	0
Djarv ⁵² 2018	CPC	Discharge/30 days	15	7	6	0	37
Duchateau ⁵³ 2017	CPC	Discharge	7	2	1	0	0
Evans ⁵ 2016	GOS	Discharge	24	6	15	3	97
Falcone ⁵⁴ 1995	Qualitative	Discharge	4	0	2	0	0
Fisher ⁵⁵ 1999	Qualitative	Discharge	0	0	0	1	0
Graesner ⁵⁶ 2011	GOS	Discharge	72	53	48	40	8
Hillman ⁵⁷ 2016	Qualitative	Variable	1	0	1	0	2
Huber-Wagner ⁵⁸ 2007	GOS	Discharge	15	13	15	6	81
Keller ³⁷ 2013	GOS	Variable	12	2	2	1	20
Kleber ²³ 2014	GOS	Discharge	1	3	10	1	0
Lawhon ²⁰ 1995	Qualitative	Discharge/3 years	0	0	1	1	0
Lin ⁵⁹ 2016	PCPC	Discharge	12	26‡	N/A	N/A	0
Love ⁶⁰ 2016	Qualitative	Discharge	4	0	3	0	0
Lundy ⁶¹ 2011	Qualitative	Discharge	36	0	89	0	0
Molina ⁶² 2008	Qualitative	Discharge	8	0	0	0	0
Mollberg ⁶³ 2011	Qualitative	Discharge	0	0	1	0	0
Moore ⁶⁴ 2011	Qualitative	Discharge	46	9	1	0	0
Moore ⁴⁴ 2016	GOS	Discharge	72	13	0	17	4
Murphy ⁶⁵ 2010	Qualitative	Discharge	16	3	7	2	0
Perron ⁶⁶ 2001	FIM	Discharge	177§	0	7	0	0
Pickens ⁶⁷ 2005	Qualitative	Discharge	11	1	0	2	0
Powell ⁶⁸ 2004	Qualitative	Discharge	15	6	5	0	36
Rabinovic ⁶⁹ 2014	Qualitative	Discharge	7	1	1	0	0
Shimazu ⁷⁰ 1983	Qualitative	30 days post admission	4	0	2	0	1
Stratton ⁷¹ 1998	CPC	Discharge	3	1	2	3	0
Tarmey ⁴³ 2011	GOS	Discharge	4	0	0	0	0
Vassallo ⁷² 2019	GOS	Discharge/30 days	2	2	1	0	2
Zwingmann ⁷³ 2015	GOS	After resuscitation	14	13	6	10	0
Zwingmann ²¹ 2016	GOS	After resuscitation	193	201	174	81	0
Total		Favorable: 992			Unfavorable: 630		Missing: 546

*Reported as good or moderate.

†Reported as poor or vegetative.

‡Reported as moderate, poor, or vegetative.

§Reported as good, moderate, or poor.

CPC, Cerebral Performance Categories; FIM, Functional Independence Measure; GOS, Glasgow Outcome Scale; N/A, not available; PCPC, Pediatric Cerebral Performance Categories.

on daily support had to be used as indicators of neurological status. Furthermore, as an outcome measurement scale was not used, there is likely some degree of variation and biases among physician-reported outcomes. Several included studies reported dichotomous outcomes using a standardized scale by combining CPC 1 to 2 (good and moderate) as a “good” outcome and

CPC 3 to 5 (severe, vegetative, death) as a “poor” outcome. However, the definition of a “good” and “poor” outcome varied between studies as CPC 3 was historically considered a “good” outcome.²⁴ This reflects the differing values and preferences in the evaluation of neurological outcomes after arrest; therefore, reporting the exact deficits of the survivors is a key component

Table 3 Quality review of included studies

Study identification	Confounding*	Participant selection†	Classification of interventions‡	Deviation from intended interventions§	Missing data¶	Outcomes**	Selective reporting††	Overall bias
Alanezi ⁴⁵ 2004	Serious	Moderate	Low	Moderate	Low	Serious	Moderate	Serious
Barnard ⁴ 2017	Serious	Moderate	Low	Moderate	Moderate	Moderate	Moderate	Serious
Beck ⁴⁶ 2016	Moderate	Moderate	Low	Low	Moderate	Moderate	Moderate	Moderate
Calkins ³⁴ 2002	Serious	Low	Low	Low	Low	Serious	Moderate	Serious
Capizzani ³⁵ 2010	Moderate	Low	Low	Low	Low	Serious	Moderate	Serious
Chia ⁴⁷ 2017	Moderate	Low	Low	Moderate	Low	Moderate	Moderate	Moderate
Chiang ⁴⁸ 2015	Moderate	Low	Low	Moderate	Low	Moderate	Low	Moderate
Chien ⁴⁹ 2016	Serious	Low	Low	Low	Moderate	Moderate	Moderate	Serious
David ⁵⁰ 2007	Moderate	Low	Low	Moderate	Low	Moderate	Moderate	Moderate
Deasy ⁵¹ 2012	Moderate	Low	Low	Low	Serious	Moderate	Moderate	Serious
Di Bartolomeo ¹⁹ 2005	Serious	Low	Low	Low	Low	Moderate	Moderate	Serious
Djarv ⁵² 2018	Moderate	Low	Low	Low	Serious	Moderate	Moderate	Serious
Duchateau ³³ 2017	Moderate	Low	Low	Low	Low	Moderate	Moderate	Moderate
Evans ⁴⁴ 2016	Moderate	Low	Low	Low	Moderate	Moderate	Moderate	Moderate
Falcone ⁵⁴ 1995	Serious	Low	Low	Moderate	Low	Serious	Moderate	Serious
Fisher ⁵⁵ 1999	Serious	Low	Low	Moderate	Low	Serious	Moderate	Serious
Graesner ⁵⁶ 2011	Moderate	Low	Low	Moderate	Moderate	Moderate	Moderate	Moderate
Hillman ⁵⁷ 2016	Serious	Moderate	Low	Moderate	Serious	Serious	Moderate	Serious
Huber-Wagner ⁵⁸ 2007	Moderate	Moderate	Low	Moderate	Serious	Moderate	Moderate	Serious
Keller ³⁷ 2013	Serious	Serious	Low	Low	Serious	Moderate	Moderate	Serious
Kleber ²³ 2014	Serious	Low	Low	Moderate	Low	Moderate	Moderate	Serious
Lawhon ²⁰ 1995	Serious	Low	Low	Low	Low	Serious	Moderate	Serious
Lin ⁵⁹ 2016	Moderate	Moderate	Low	Low	Low	Moderate	Moderate	Moderate
Love ⁶⁰ 2016	Moderate	Low	Low	Moderate	Low	Serious	Moderate	Serious
Lundy ⁶¹ 2011	Moderate	Moderate	Low	Moderate	Low	Moderate	Moderate	Moderate
Molina ⁶² 2008	Serious	Low	Low	Low	Low	Serious	Moderate	Serious
Mollberg ⁶³ 2011	Serious	Low	Low	Moderate	Low	Serious	Moderate	Serious
Moore ⁶⁴ 2011	Moderate	Low	Low	Low	Low	Moderate	Moderate	Serious
Moore ⁴⁴ 2016	Moderate	Low	Low	Low	Low	Moderate	Moderate	Moderate
Murphy ⁶⁵ 2010	Serious	Low	Low	Low	Low	Serious	Moderate	Serious
Perron ⁶⁶ 2001	Moderate	Low	Low	Low	Serious	Moderate	Serious	Serious
Pickens ⁶⁷ 2005	Moderate	Moderate	Low	Low	Low	Moderate	Moderate	Moderate
Powell ⁶⁸ 2004	Serious	Serious	Low	Low	Low	Moderate	Moderate	Serious
Rabinovici ⁶⁹ 2014	Moderate	Low	Low	Low	Low	Serious	Moderate	Serious
Shimazu ⁷⁰ 1983	Serious	Moderate	Moderate	Low	Low	Serious	Moderate	Serious
Stratton ⁷¹ 1998	Moderate	Low	Low	Low	Low	Moderate	Moderate	Moderate
Tarmey ⁴³ 2011	Serious	Low	Low	Moderate	Low	Moderate	Moderate	Serious
Vassallo ⁷² 2019	Serious	Moderate	Low	Low	Moderate	Moderate	Moderate	Serious
Zwingmann ⁷³ 2015	Moderate	Serious	Low	Low	Low	Moderate	Moderate	Serious
Zwingmann ²¹ 2016	Moderate	Serious	Moderate	Moderate	Low	Moderate	Moderate	Serious

*Were confounding variables controlled for with statistical analysis (eg, multivariate analysis)? Did they determine these factors appropriately (eg, univariate analysis, references, subject matter expert, etc)?

†Were only survivors/non-vegetative participants assessed? Was any group excluded (eg, specific year of arrests, no rhythm on presentation, type of trauma, witnessed cardiac arrest only, etc)?

‡Was the exposure (traumatic cardiac arrest) well defined before the study? Was there a method used to identify these cases?

§Were special methods used for resuscitation (eg, therapeutic hypothermia, thoracotomy)? If so, was it reported and appropriately accounted for?

¶Were any patients lost to follow-up? Was any method used to account for this/any attempt made to recover from this? Was there a particular group of patients that were lost to follow-up?

**Was the exposure known to the person assessing neurological outcome? How were neurological outcomes assessed? Was it subjective/unclear (eg, physician-reported, self-reported, etc)? Was a standardized scale used and data reported for all categories (eg, Glasgow Outcome Scale)?

††Was there a preregistered protocol with preplanned outcomes to report? Were only specific outcomes reported in full? Were the neurological outcomes dichotomized into "good" and "poor" instead of reporting for each category?

of their neuroprognostication, as there may be a large difference between a moderate disability and a fully recovered patient.

Although the GOS and CPC scales were commonly used, alternatives such as the mRS and the GOSE were not used in any of the included studies. The CPC is heavily weighted toward mental functions and has been criticized for being inadequate to assess functional status at hospital discharge,^{25 26} which is supported by significant variability in quality-of-life measures for patients with similar CPC scores.^{27 28} Alternatively, the mRS and GOSE scales consider work capacities, social activities, and return to social life.^{9 11 29} However, most studies assessed neurological status at discharge, which may be inadequate regardless of the scale used. Most criteria rely on whether patients can perform daily activities, which are not undertaken while in hospital.¹¹ As cardiac arrest survivors tend to report cognitive impairment and restricted societal participation after hospital discharge,³⁰ early assessments may overestimate the neurological outcomes. Hence, repeated assessments over time with multiple scales are recommended to accurately assess the functional progression of these patients.

This review has several limitations. To start, our overall neurological outcome may be an overestimate given that patients with moderate deficits and favorable discharge dispositions were assumed to have a favorable outcome. Furthermore, the favorable neurological recovery rate that we provide here is based on a diverse patient population and does not account for the mechanism of trauma (blunt vs. penetrating), the patient age group (adult vs. pediatric), the time point when neurological status is assessed, or special resuscitative techniques. Previous literature has demonstrated varying effects of these factors in neurological outcomes. The effect of the mechanism of injury on traumatic arrest outcomes has been controversial. Whereas some cohort studies suggest that good outcomes can be achieved in select patient groups, especially in those with penetrating injuries,^{31 32} other studies suggest that the mechanism of injury is not associated with better outcomes (neurological recovery or survival).^{22 33} Hence, it would be valuable to assess the effect of the mechanism of injury on the neurological recovery following arrest. Although only a few studies assess traumatic arrest in the pediatric population, there is a general consensus that the functional outcomes are poorer compared with adults.^{15 34 35} In the length of follow-up after cardiac arrest, there is evidence that significant recovery can occur between 1 and 6 months³⁶ and long-term recovery after 1 year.³⁷ The majority of the studies we included assessed neurological outcomes at patient discharge or after resuscitation (table 2), suggesting that we underestimated the prevalence of good neurological recovery in the current review. Special resuscitative techniques, such as emergency thoracotomy and therapeutic hypothermia, have some demonstrated benefits in neurological outcomes after traumatic arrest for select patient populations.^{38–42} Based on the effects of these factors on neurological recovery, we recommend for future studies evaluating outcomes after traumatic arrest (1) reporting the mechanism of injury for survivors in each neurological outcome category, (2) reporting the age group (adult vs. pediatric) in each neurological outcome category, (3) assessing neurological outcomes 1 year after the arrest if resources allow or >30 days if necessary,³⁶ and (4) reporting special resuscitative techniques in individual survivors in each neurological outcome category.

Overall, the strength of the scientific evidence for neurological outcomes after TCA is low. Although we identified some large national registry studies, most studies were retrospective cohorts.^{4 5 43} Five prospective studies were included, but only one was a multicenter study.⁴⁴ Furthermore, our quality assessment

revealed a moderate to serious risk of bias in our included studies due to confounding and selective reporting domains. Our outcome of interest was often qualitatively described, and some studies briefly reported neurological outcomes for only one category (eg, vegetative state). As a result, many studies without sufficient data were excluded, and the extent of the selective reporting bias we identified may be an underestimate.

Good and moderate neurological outcomes are frequently reported in patients who survive TCA. However, stronger evidence is needed to prognosticate this patient population as neurological outcomes are often inadequately reported. Future studies should identify and adjust for appropriate confounding variables and report the prevalence of each neurological category. Multicenter prospective studies that focus on the quality of survivorship are urgently needed.

Acknowledgements We thank Si Hong Yu (Health Sciences Librarian at Queen's University) for his help with the development of the search strategy.

Contributors DS, CM, and CE conceived the study design. DS and CM created the literature search. DS and CM collected data and conducted the quality assessment with assistance from CE. DS drafted the article. CM prepared the figure and tables, and all authors contributed substantially to revision. All authors take responsibility for the article as a whole. DS is the guarantor of the article and accepts responsibility for the integrity of the published work.

Funding External funding was received from the Queen's Department of Emergency Medicine in the Emergency Medicine Summer Studentship Award (DS and CM).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Kauvar DS, Lefering R, Wade CE. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma* 2006;60:S3–11.
- 2 Millin MG, Galvagno SM, Khandker SR, Malki A, Bulger EM, . Standards and Clinical Practice Committee of the National Association of EMS Physicians (NAEMSP) Subcommittee on Emergency Services—Prehospital of the American College of Surgeons' Committee on Trauma (ACSCOT). Withholding and termination of resuscitation of adult cardiopulmonary arrest secondary to trauma: resource document to the joint NAEMSP-ACSCOT position statements. *J Trauma Acute Care Surg* 2013;75:459–67.
- 3 Rosemurgy AS, Norris PA, Olson SM, Hurst JM, Albrink MH. Prehospital traumatic cardiac arrest: the cost of futility. *J Trauma* 1993;35:468–72.
- 4 Barnard E, Yates D, Edwards A, Fragoso-Iñiguez M, Jenks T, Smith JE. Epidemiology and aetiology of traumatic cardiac arrest in England and Wales - A retrospective database analysis. *Resuscitation* 2017;110:90–4.
- 5 Evans CCD, Petersen A, Meier EN, Buick JE, Schreiber M, Kannas D, Austin MA, . Resuscitation Outcomes Consortium Investigators. Prehospital traumatic cardiac arrest: management and outcomes from the resuscitation outcomes Consortium epistry-trauma and PROPHET registries. *J Trauma Acute Care Surg* 2016;81:285–93.
- 6 Leis CC, Hernández CC, Blanco MJG-O, Paterna PCR, Hernández RdeE, Torres EC. Traumatic cardiac arrest: should advanced life support be initiated? *J Trauma Acute Care Surg* 2013;74:634–8.
- 7 Smith JE, Rickard A, Wise D. Traumatic cardiac arrest. *J R Soc Med* 2015;108:11–16.
- 8 Kim Y-J, Ahn S, Sohn CH, Seo D-W, Lee Y-S, Lee JH, Oh BJ, Lim KS, Kim WY. Long-Term neurological outcomes in patients after out-of-hospital cardiac arrest. *Resuscitation* 2016;101:1–5.
- 9 Sandroni C, D'Arrigo S, Nolan JP. Prognostication after cardiac arrest. *Crit Care* 2018;22:150.
- 10 Hofgren C, Lundgren-Nilsson A, Esbjörnsson E, Sunnerhagen KS. Two years after cardiac arrest; cognitive status, ADL function and living situation. *Brain Inj* 2008;22:972–8.
- 11 Rittenberger JC, Raina K, Holm MB, Kim YJ, Callaway CW. Association between cerebral performance category, modified Rankin scale, and discharge disposition after cardiac arrest. *Resuscitation* 2011;82:1036–40.

- 12 Weir J, Steyerberg EW, Butcher I, Lu J, Lingsma HF, McHugh GS, Rozenbeek B, Maas AIR, Murray GD. Does the extended Glasgow outcome scale add value to the conventional Glasgow outcome scale? *J Neurotrauma* 2012;29:53–8.
- 13 Battistella FD, Nugent W, Owings JT, Anderson JT. Field triage of the pulseless trauma patient. *Arch Surg* 1999;134:742.
- 14 Martin SK, Shatney CH, Sherck JP, Ho C-C, Homan SJ, Neff J, Moore EE. Blunt trauma patients with prehospital pulseless electrical activity (pea): poor ending assured. *J Trauma* 2002;53:876–81.
- 15 Zwingmann J, Mehlhorn AT, Hammer T, Bayer J, Südkamp NP, Strohm PC. Survival and neurologic outcome after traumatic out-of-hospital cardiopulmonary arrest in a pediatric and adult population: a systematic review. *Crit Care* 2012;16:R117.
- 16 Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JAC, et al. The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- 17 Veritas Health Innovation. Covidence systematic review software. Melbourne, Australia: Veritas Health Innovation, 2019.
- 18 McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med* 2012;22:276–82.
- 19 Di Bartolomeo S, Sanson G, Nardi G, Michelutto V, Scian F. Hems vs. Ground-BLS care in traumatic cardiac arrest. *Prehosp Emerg Care* 2005;9:79–84.
- 20 Lawhon JC, Somberg LB, Barker DE. Prehospital blunt traumatic arrest: survival and analysis. *J Tenn Med Assoc* 1995;88:4.
- 21 Zwingmann J, Lefering R, Feucht M, Südkamp NP, Strohm PC, Hammer T. Outcome and predictors for successful resuscitation in the emergency room of adult patients in traumatic cardiorespiratory arrest. *Crit Care* 2016;20:282.
- 22 Kim JG, Lee J, Choi HY, Kim W, Kim J, Moon S, Shin H, Ahn C, Cho Y, Shin DG, et al. Outcome analysis of traumatic out-of-hospital cardiac arrest patients according to the mechanism of injury: a nationwide observation study. *Medicine* 2020;99:e23095.
- 23 Kleber C, Giesecke MT, Lindner T, Haas NP, Buschmann CT. Requirement for a structured algorithm in cardiac arrest following major trauma: epidemiology, management errors, and preventability of traumatic deaths in Berlin. *Resuscitation* 2014;85:405–10.
- 24 Sandroni C, Nolan JP. Neuroprognostication after cardiac arrest in Europe: new timings and standards. *Resuscitation* 2015;90:A4–5.
- 25 Cummins RO, Chamberlain D, Hazinski MF, Nadkarni V, Kloeck W, Kramer E, Becker L, Robertson C, Koster R, Zaritsky A, et al. Recommended guidelines for reviewing, reporting, and conducting research on in-hospital resuscitation: the in-hospital "Utstein style". American Heart Association. *Ann Emerg Med* 1997;29:650–79.
- 26 Hsu* JWY, Madsen§ CD, Callahan ML. Quality-Of-Life and formal functional testing of survivors of out-of-hospital cardiac arrest correlates poorly with traditional neurologic outcome scales. *Ann Emerg Med* 1996;28:597–605.
- 27 Raina KD, Callaway C, Rittenberger JC, Holm MB. Neurological and functional status following cardiac arrest: method and tool utility. *Resuscitation* 2008;79:249–56.
- 28 Stiell IG, Nesbitt LP, Nichol G, Maloney J, Dreyer J, Beaudoin T, Blackburn J, Wells GA, OPALS Study Group. Comparison of the cerebral performance category score and the health Utilities index for survivors of cardiac arrest. *Ann Emerg Med* 2009;53:241–8.
- 29 Teasdale GM, Pettigrew LE, Wilson JT, Murray G, Jennett B. Analyzing outcome of treatment of severe head injury: a review and update on advancing the use of the Glasgow outcome scale. *J Neurotrauma* 1998;15:587–97.
- 30 Haywood K, Whitehead L, Nadkarni VM, Achana F, Beesems S, Böttiger BW, Brooks A, Castrén M, Ong ME, Hazinski MF, et al. COSCA (core outcome set for cardiac arrest) in adults: an Advisory statement from the International liaison Committee on resuscitation. *Circulation* 2018;137:e783–801.
- 31 Dane FC, Russell-Lindgren KS, Parish DC, Durham MD, Brown TD. In-Hospital resuscitation: association between ACLS training and survival to discharge. *Resuscitation* 2000;47:83–7.
- 32 Davies GE, Lockett DJ. Thirteen survivors of prehospital thoracotomy for penetrating trauma: a prehospital physician-performed resuscitation procedure that can yield good results. *J Trauma* 2011;70:E75–8.
- 33 Tran A, Fernando SM, Rochweg B, Vaillancourt C, Inaba K, Kyremanteng K, Nolan JP, McCredie VA, Petrosiak A, Hicks C, et al. Pre-arrest and intra-arrest prognostic factors associated with survival following traumatic out-of-hospital cardiac arrest - A systematic review and meta-analysis. *Resuscitation* 2020;153:119–35.
- 34 Calkins CM, Bensard DD, Partrick DA, Karrer FM. A critical analysis of outcome for children sustaining cardiac arrest after blunt trauma. *J Pediatr Surg* 2002;37:180–4.
- 35 Capizzani AR, Drongowski R, Ehrlich PF. Assessment of termination of trauma resuscitation guidelines: are children small adults? *J Pediatr Surg* 2010;45:903–7.
- 36 Geocadin RG, Callaway CW, Fink EL, Golan E, Greer DM, Ko NU, Lang E, Licht DJ, Marino BS, McNair ND, et al. Standards for studies of neurological prognostication in comatose survivors of cardiac arrest: a scientific statement from the American heart association. *Circulation* 2019;140:e517–42.
- 37 Keller D, Kulp H, Maher Z, Santora TA, Goldberg AJ, Seamon MJ. Life after near death: long-term outcomes of emergency department thoracotomy survivors. *J Trauma Acute Care Surg* 2013;74:1315–20.
- 38 Baker CC, Caronna JJ, Trunkey DD. Neurologic outcome after emergency room thoracotomy for trauma. *Am J Surg* 1980;139:677–81.
- 39 Hughes M, Perkins Z. Outcomes following resuscitative thoracotomy for abdominal exsanguination, a systematic review. *Scand J Trauma Resusc Emerg Med* 2020;28:9.
- 40 Seamon MJ, Haut ER, Van Arendonk K, Barbosa RR, Chiu WC, Dente CJ, Fox N, Jawa RS, Khwaja K, Lee JK, et al. An evidence-based approach to patient selection for emergency department thoracotomy: a practice management guideline from the eastern association for the surgery of trauma. *J Trauma Acute Care Surg* 2015;79:159–73.
- 41 Song SS, Lyden PD. Overview of therapeutic hypothermia. *Curr Treat Options Neurol* 2012;14:541–8.
- 42 Tuma MA, Stansbury LG, Stein DM, McQuillan KA, Scalea TM. Induced hypothermia after cardiac arrest in trauma patients: a case series. *J Trauma* 2011;71:1524–7.
- 43 Tarmey NT, Park CL, Bartels OJ, Konig TC, Mahoney PF, Mellor AJ. Outcomes following military traumatic cardiorespiratory arrest: a prospective observational study. *Resuscitation* 2011;82:1194–7.
- 44 Moore EE, Knudson MM, Burlew CC, Inaba K, Dicker RA, Biffi WL, Malhotra AK, Schreiber MA, Browder TD, Coimbra R, et al. Defining the limits of resuscitative emergency department thoracotomy: a contemporary Western trauma association perspective. *J Trauma* 2011;70:334–9.
- 45 Alanezi K, Alanzi F, Faidi S, Sprague S, Cadeddu M, Baillie F, Bowser D, McCallum A, Bhandari M. Survival rates for adult trauma patients who require cardiopulmonary resuscitation. *CJEM* 2004;6:263–5.
- 46 Beck B, Tohira H, Bray JE, Straney L, Brown E, Inoue M, Williams TA, McKenzie N, Celena A, Bailey P, et al. Trends in traumatic out-of-hospital cardiac arrest in Perth, Western Australia from 1997 to 2014. *Resuscitation* 2016;98:79–84.
- 47 Chia MY-C, Lu QS, Rahman NH, Doctor NE, Nishiuchi T, Leong BS-H, Tham LP, Goh E-S, Tiah L, Monsomboon A, et al. Characteristics and outcomes of young adults who suffered an out-of-hospital cardiac arrest (OHCA). *Resuscitation* 2017;111:34–40.
- 48 Chiang W-C, Chen S-Y, Ko PC-I, Hsieh M-J, Wang H-C, Huang EP-C, Yang C-W, Chong K-M, Chen W-T, Chen S-Y, et al. Prehospital intravenous epinephrine may boost survival of patients with traumatic cardiac arrest: a retrospective cohort study. *Scand J Trauma Resusc Emerg Med* 2015;23:102.
- 49 Chien C-Y, Su Y-C, Lin C-C, Kuo C-W, Lin S-C, Weng Y-M. Is 15 minutes an appropriate resuscitation duration before termination of a traumatic cardiac arrest? A case-control study. *Am J Emerg Med* 2016;34:505–9.
- 50 David J-S, Gueugniaud P-Y, Riou B, Pham E, Dubien P-Y, Goldstein P, Freys M, Petit P, Pierre-Yves MD, Riou Ph D, Bruno MD, Pham Ph D, PhD E. Does the prognosis of cardiac arrest differ in trauma patients? *Crit Care Med* 2007;35:2251–5.
- 51 Deasy C, Bray J, Smith K, Harris L, Morrison C, Bernard S, Cameron P. Traumatic out-of-hospital cardiac arrests in Melbourne, Australia. *Resuscitation* 2012;83:465–70.
- 52 Djarv T, Axelsson C, Herlitz J, Stromsoe A, Israelsson J, Claesson A. Traumatic cardiac arrest in Sweden 1990-2016 - a population-based national cohort study. *Scand J Trauma Resusc Emerg Med* 2018;26:30.
- 53 Duchateau F-X, Hamada S, Raux M, Gay M, Mantz J, Paugam Burtz C, Gauss T, Traumabase Group. Long-Term prognosis after out-of-hospital resuscitation of cardiac arrest in trauma patients: prehospital trauma-associated cardiac arrest. *Emerg Med J* 2017;34:34–8.
- 54 Falcone RE, Herron H, Johnson R, Childress S, Lacey P, Scheiderer G. Air medical transport for the trauma patient requiring cardiopulmonary resuscitation: a 10-year experience. *Air Med J* 1995;14:197–204.
- 55 Fisher B, Worthen M. Cardiac arrest induced by blunt trauma in children. *Pediatr Emerg Care* 1999;15:274–6.
- 56 Gräsner JT, Wnent J, Seewald S, Meybohm P, Fischer M, Paffrath T, Wafaisade A, Bein B, Lefering R. German resuscitation registry Working group, trauma registry of the German Society for trauma surgery (DGU). cardiopulmonary resuscitation traumatic cardiac arrest-there are survivors. An analysis of two national emergency registries. *Crit Care* 2011;15:R276.
- 57 Hillman CM, Rickard A, Rawlins M, Smith JE. Paediatric traumatic cardiac arrest: data from the joint theatre trauma registry. *J R Army Med Corps* 2016;162:276–9.
- 58 Huber-Wagner S, Lefering R, Quick M, Kay MV, Paffrath T, Mutschler W, Kanz KG. Working group on Polytrauma of the German Trauma Society (DGU). outcome in 757 severely injured patients with traumatic cardiorespiratory arrest. *Resuscitation* 2007;75:276–85.
- 59 Lin Y-R, Syue Y-J, Buddhakosai W, Lu H-E, Chang C-F, Chang C-Y, Chen CH, Chen W-L, Li C-J, HE L, Chang CY, CJ L. Impact of different initial epinephrine treatment time points on the early Postresuscitative hemodynamic status of children with traumatic out-of-hospital cardiac arrest. *Medicine* 2016;95:e3195.
- 60 Love KM, Brown JB, Harbrecht BG, Muldoon SB, Miller KR, Bennis MV, Smith JW, Baker CE, Franklin GA. Organ donation as an outcome of traumatic cardiopulmonary arrest: a cost evaluation. *J Trauma Acute Care Surg* 2016;80:792–8.
- 61 Lundy DJ, Ross SE, Schorr C, Jones AE, Trzeciak S. Outcomes of trauma victims with cardiac arrest who survive to intensive care unit admission. *J Trauma* 2011;71:E12–16.
- 62 Molina EJ, Gaughan JP, Kulp H, McClurken JB, Goldberg AJ, Seamon MJ. Outcomes after emergency department thoracotomy for penetrating cardiac injuries: a new perspective. *Interact Cardiovasc Thorac Surg* 2008;7:845–8.
- 63 Mollberg NM, Wise SR, Berman K, Chowdhry S, Holevar M, Sullivan R, Vafa A. The consequences of noncompliance with guidelines for withholding or terminating resuscitation in traumatic cardiac arrest patients. *J Trauma* 2011;71:997–1002.
- 64 Moore HB, Moore EE, Burlew CC, Biffi WL, Pieracci FM, Barnett CC, Bensard DD, Jurkovich GJ, Fox CJ, Sauaia A. Establishing benchmarks for resuscitation of traumatic

- circulatory arrest: Success-to-Rescue and survival among 1,708 patients. *J Am Coll Surg* 2016;223:42–50.
- 65 Murphy JT, Jaiswal K, Sabella J, Vinson L, Megison S, Maxson RT. Prehospital cardiopulmonary resuscitation in the pediatric trauma patient. *J Pediatr Surg* 2010;45:1413–9.
- 66 Perron AD, Sing RF, Branas CC, Huynh T. Predicting survival in pediatric trauma patients receiving cardiopulmonary resuscitation in the prehospital setting. *Prehosp Emerg Care* 2001;5:6–9.
- 67 Pickens JJ, Copass MK, Bulger EM. Trauma patients receiving CPR: predictors of survival. *J Trauma* 2005;58:951–8.
- 68 Powell DW, Moore EE, Cothren CC, Ciesla DJ, Burch JM, Moore JB, Johnson JL. Is emergency department resuscitative thoracotomy futile care for the critically injured patient requiring prehospital cardiopulmonary resuscitation? *J Am Coll Surg* 2004;199:211–5.
- 69 Rabinovici R. Sixty-Seven consecutive resuscitative Thoracotomies by a single surgeon. *Scand J Surg* 2014;103:156–60.
- 70 Shimazu S, Shatney CH. Outcomes of trauma patients with no vital signs on hospital admission. *J Trauma* 1983;23:213–6.
- 71 Stratton SJ, Brickett K, Crammer T. Prehospital pulseless, unconscious penetrating trauma victims: field assessments associated with survival. *J Trauma* 1998;45:96–100.
- 72 Vassallo J, Webster M, Barnard EBG, Lyttle MD, PERUKI Smith JE;. Paediatric emergency research in the UK and Ireland). epidemiology and aetiology of paediatric traumatic cardiac arrest in England and Wales. *Arch Dis Child* 2019;104:437–43.
- 73 Zwingmann J, Lefering R, Bayer J, Reising K, Kuminack K, Südkamp NP, Strohm PC. TraumaRegister DGU(®). Outcome and risk factors in children after traumatic cardiac arrest and successful resuscitation. *Resuscitation* 2015;96:59–65.